arXiv:1504.07813v1 [math.QA] 29 Apr 2015

Explicit Forms of Cluster Variables on Double Bruhat Cells $G^{u,e}$ of type C

YUKI KANAKUBO* and TOSHIKI NAKASHIMA†

Dedicated to Professor Ken-ichi SHINODA

Abstract

Let $G = Sp_{2r}(\mathbb{C})$ be a simply connected simple algebraic group over $\mathbb{C}$ of type C_r, B and B_- be its two opposite Borel subgroups, and W be the associated Weyl group. For u, $v \in W$, it is known that the coordinate ring $\mathbb{C}[G^{u,v}]$ of the double Bruhat cell $G^{u,v} = BuB \cup B_-vB_-$ is isomorphic to an upper cluster algebra $\overline{\mathcal{A}}(\mathbf{i})_{\mathbb{C}}$ and the generalized minors $\Delta(k; \mathbf{i})$ are the cluster variables of $\mathbb{C}[G^{u,v}]$[5]. In the case $v = e$, we shall describe the generalized minor $\Delta(k; \mathbf{i})$ explicitly.

Contents

1 Introduction 2

2 Fundamental representations for type C_r 3

3 Factorization theorem for type C 4
- 3.1 Double Bruhat cells 4
- 3.2 Factorization theorem for type C_r 5

4 Cluster algebras and generalized minors 8
- 4.1 Cluster algebras of geometric type 8
- 4.2 Upper cluster algebra $\overline{\mathcal{A}}(\mathbf{i})$ 9
- 4.3 Generalized minors and bilinear form 10
- 4.4 Cluster algebras on Double Bruhat cells of type C 11

5 Generalized minors and crystals 11
- 5.1 Generalized minor $\Delta^G(k; \mathbf{i})(a, \mathbf{Y})$ 12
- 5.2 Generalized minor $\Delta^L(k; \mathbf{i})(\mathbf{Y})$ 13

6 The proof of Theorem 5.7 15
- 6.1 The set $X_d(m, m')$ of paths: path descriptions 15
- 6.2 One-to-one correspondence between paths in $X_d(m, m')$ and terms of $\Delta^L(k; \mathbf{i})(\mathbf{Y})$ 19
- 6.3 The properties of paths in $X_d(m, m')$ 23
- 6.4 The proof of Theorem 5.7 27

---

*Division of Mathematics, Sophia University, Kioicho 7-1, Chiyoda-ku, Tokyo 102-8554, Japan: j_chi_sen_you_ky@sophia.ac.jp

†Division of Mathematics, Sophia University, Kioicho 7-1, Chiyoda-ku, Tokyo 102-8554, Japan: toshiki@sophia.ac.jp: supported in part by JSPS Grants in Aid for Scientific Research ♯22540031.

1 Introduction

Let G be a simply connected simple algebraic group over $\mathbb{C}$ of rank r, $B, B_- \subset G$ the opposite Borel subgroups, $H := B \cap B^-$ the maximal torus, $N \subset B$, $N_- \subset B_-$ the maximal unipotent subgroups and W the associated Weyl group. For $u, v \in W$, define $G^{u,v} := (BuB) \cap (B_-vB_-)$ (resp. $L^{u,v} := (NuN) \cap (B_-vB_-)$) and call it the (reduced) double Bruhat cell.

In [5], it is shown that the coordinate ring $\mathbb{C}[G^{u,v}]$ $(u, v \in W)$ of double Bruhat cell $G^{u,v}$ has the structure of an upper cluster algebra. The initial cluster variables of this upper cluster algebras are given as certain generalized minors on $G^{u,v}$.

In [7], we treated the case of type A and $v = e$, where we described the explicit forms of the generalized minors $\{\Delta(k; \mathbf{i})\}$ and revealed the linkage between $\Delta(k; \mathbf{i})$ and monomial realizations of crystals.

In this paper, we shall write down the explicit forms of the generalized minors $\Delta(k; \mathbf{i})$ on the (reduced) double Bruhat cell $G^{u,e}$ ($L^{u,e}$) of type C_r by using the 'path descriptions' of generalized minors (see Sect.6), where we only treat a Weyl group elements u with the form as in (3.2) and denote its reduced word $\mathbf{i}$ by (3.3). Indeed, generalized minors are expressed in terms of certain invariant bilinear forms (see (4.3)). And then, using this bilinearity we obtain 'path descriptions' of the generalized minors.

Unfortunately, we do not present the relation between the explicit forms of $\Delta(k; \mathbf{i})$ and crystals here unlike with [7]. We will, however, discuss this elsewhere.

The main result is given as in Theorem 5.7: Let $\mathbf{i}$ be the reduced word of u as above and i_k is the k-th index of $\mathbf{i}$ form the left. In [2], it is shown that there exists a biregular isomorphism from $(\mathbb{C}^\times)^n$ to a Zariski open subset of $L^{u,e}$ $(n := l(u))$ (see Theorem 3.3). We denote this isomorphism by $x^L_{\mathbf{i}}$ and set $\Delta^L(k; \mathbf{i}) := \Delta(k; \mathbf{i}) \circ x^L_{\mathbf{i}}$. We also set the monomials $\overline{C}(l, k)$ and $C(l, k)$ as in (5.11).
Theorem 5.7 We set $d := i_k = i_n$ and

$$\mathbf{Y} := (Y_{1,1}, Y_{1,2}, \cdots, Y_{1,r}, \cdots, Y_{m-1,1}, \cdots, Y_{m-1,r}, Y_{m,1}, \cdots, Y_{m,i_n}) \in (\mathbb{C}^\times)^n.$$

Then we have

$$\begin{aligned}\Delta^L(k; \mathbf{i})(\mathbf{Y}) = \sum_{(*)} \prod_{i=1}^{d} &\overline{C}(m - l_i^{(1)}, k_i^{(1)}) \cdots \overline{C}(m - l_i^{(\delta_i)}, k_i^{(\delta_i)}) \\ &\cdot C(m - l_i^{(\delta_i+1)}, |k_i^{(\delta_i+1)}| - 1) \cdots C(m - l_i^{(m-m')}, |k_i^{(m-m')}| - 1),\end{aligned}$$

$$l_i^{(s)} := \begin{cases} k_i^{(s)} + s - i - 1 & \text{if } s \le \delta_i, \\ s - i + r & \text{if } s > \delta_i \end{cases} \qquad (1 \le i \le d)$$

where $(*)$ is the conditions for $k_i^{(s)}$ $(1 \le s \le m - m',\ 1 \le i \le d)$: $1 \le k_1^{(s)} < k_2^{(s)} < \cdots < k_d^{(s)} \le \overline{1}$ $\ (1 \le s \le m - m')$, $1 \le k_i^{(1)} \le \cdots \le k_i^{(m-m')} \le m' + i$ $\ (1 \le i \le r - m')$, and $1 \le k_i^{(1)} \le \cdots \le k_i^{(m-m')} \le \overline{1}$ $\ (r - m' + 1 \le i \le d)$, and δ_i $(i = 1, \cdots, d)$ are the numbers which satisfy $1 \le k_i^{(1)} \le k_i^{(2)} \le \cdots \le k_i^{(\delta_i)} \le r$, $\overline{r} \le k_i^{(\delta_i+1)} \le \cdots \le k_i^{(m-m')} \le \overline{1}$.

For $\mathbf{k}=(k_i^{(s)})$ and $\mathbf{k}'=(k_i^{'(s)})$ satisfying $(*)$, let us write the monomial

$$C(\mathbf{k}):=\prod_{i=1}^{d}\overline{C}(m-l_i^{(1)},k_i^{(1)})\cdots\overline{C}(m-l_i^{(\delta_i)},k_i^{(\delta_i)})$$
$$\cdot C(m-l_i^{(\delta_i+1)},|k_i^{(\delta_i+1)}|-1)\cdots C(m-l_i^{(m-m')},|k_i^{(m-m')}|-1).$$

Note that even if $\mathbf{k}\neq\mathbf{k}'$, we may have $C(\mathbf{k})=\mathbf{C}(\mathbf{k}')$. Thus, we will know that the coefficients of the monomials in $\Delta^L(k;\mathbf{i})$ are not necessarily 1 (See Example 5.8). We shall show Theorem 5.7 in the last section by using "path descriptions". By this theorem, we find that all the generalized minors $\{\Delta^L(k;\mathbf{i})(\mathbf{Y})\}$ are Laurent polynomials with non-negative coefficients.

Finally, we also define $\Delta^G(k;\mathbf{i}):=\Delta(k;\mathbf{i})\circ\overline{x}^G_{\mathbf{i}}$, where $\overline{x}^G_{\mathbf{i}}$ is a biregular isomorphism from $H\times(\mathbb{C}^\times)^n$ to a Zariski open subset of $G^{u,e}$ (see Proposition 3.4). In Proposition 5.3, we shall show that $\Delta^G(k;\mathbf{i})$ is immediately obtained from $\Delta^L(k;\mathbf{i})$.

2 Fundamental representations for type C_r

Let $I:=\{1,\cdots,r\}$ be a finite index set and $A=(a_{ij})_{i,j\in I}$ be the Cartan matrix of type C_r: That is, $A=(a_{i,j})_{i,j\in I}$ is given by

$$a_{i,j}=\begin{cases}2 & \text{if } i=j,\\ -1 & \text{if } |i-j|=1 \text{ and } (i,j)\neq(r-1,r),\\ -2 & \text{if } (i,j)=(r-1,r),\\ 0 & \text{otherwise.}\end{cases}$$

Let $(\mathfrak{h},\{\alpha_i\}_{i\in I},\{h_i\}_{i\in I})$ be the associated root data satisfying $\alpha_j(h_i)=a_{ij}$ where $\alpha_i\in\mathfrak{h}^*$ is a simple root and $h_i\in\mathfrak{h}$ is a simple co-root. Note that α_i $(i\neq r)$ are short roots and α_r is the long root. Let $\{\Lambda_i\}_{i\in I}$ be the set of the fundamental weights satisfying $\alpha_j(h_i)=a_{i,j}$ and $\Lambda_i(h_j)=\delta_{i,j}$. Let $P=\bigoplus_{i\in I}\mathbb{Z}\Lambda_i$ be the weight lattice and $P^*=\bigoplus_{i\in I}\mathbb{Z}h_i$ be the dual weight lattice. Define the order on the set $J:=\{i,\overline{i}|1\leq i\leq r\}$ by

$$1<2<\cdots<r-1<r<\overline{r}<\overline{r-1}<\cdots<\overline{2}<\overline{1}. \tag{2.1}$$

For the simple Lie algebra $\mathfrak{g}=\mathfrak{sp}(2r,\mathbb{C})=\langle\mathfrak{h},e_i,f_i(i\in I)\rangle$, let us describe the vector representation $V(\Lambda_1)$. Set $\mathbf{B}^{(r)}:=\{v_i,v_{\overline{i}}|i=1,2,\cdots,r\}$ and define $V(\Lambda_1):=\bigoplus_{v\in\mathbf{B}^{(r)}}\mathbb{C}v$. The weights of v_i, $v_{\overline{i}}$ $(i=1,\cdots,r)$ are as follows:

$$\mathrm{wt}(v_i)=\Lambda_i-\Lambda_{i-1},\quad \mathrm{wt}(v_{\overline{i}})=\Lambda_{i-1}-\Lambda_i, \tag{2.2}$$

where $\Lambda_0=0$. We define the $\mathfrak{sp}(2r,\mathbb{C})$-action on $V(\Lambda_1)$ as follows:

$$hv_j=\langle h,\mathrm{wt}(v_j)\rangle v_j\ \ (h\in P^*,\ j\in J), \tag{2.3}$$
$$f_iv_i=v_{i+1},\ f_iv_{\overline{i+1}}=v_{\overline{i}},\quad e_iv_{i+1}=v_i,\ e_iv_{\overline{i}}=v_{\overline{i+1}}\quad(1\leq i<r), \tag{2.4}$$
$$f_rv_r=v_{\overline{r}},\qquad e_rv_{\overline{r}}=v_r, \tag{2.5}$$

and the other actions are trivial.

Let Λ_i be the i-th fundamental weight of type C_r. As is well-known that the fundamental representation $V(\Lambda_i)$ $(1 \leq i \leq r)$ is embedded in $\wedge^i V(\Lambda_1)$ with multiplicity free. The explicit form of the highest (resp. lowest) weight vector u_{Λ_i} (resp. v_{Λ_i}) of $V(\Lambda_i)$ is realized in $\wedge^i V(\Lambda_1)$ as follows:

$$\begin{aligned} u_{\Lambda_i} &= v_1 \wedge v_2 \wedge \cdots \wedge v_i, \\ v_{\Lambda_i} &= v_{\overline{1}} \wedge v_{\overline{2}} \wedge \cdots \wedge v_{\overline{i}}. \end{aligned} \tag{2.6}$$

3 Factorization theorem for type C

In this section, we shall introduce (reduced) double Bruhat cells $G^{u,v}$, $L^{u,v}$, and their properties in the case $G = Sp_{2r}(\mathbb{C})$, $v = e$ and some special $u \in W$. In [2] and [3], these properties had been proven for simply connected, connected, semisimple complex algebraic groups and arbitrary $u, v \in W$.

For $l \in \mathbb{Z}_{>0}$, we set $[1, l] := \{1, 2, \cdots, l\}$.

3.1 Double Bruhat cells

Let $G = Sp_{2r}(\mathbb{C})$ be the simple complex algebraic group of type C_r, B and B_- be two opposite Borel subgroups in G, $N \subset B$ and $N_- \subset B_-$ be their unipotent radicals, $H := B \cap B_-$ a maximal torus. Let $W := \langle s_i | i = 1, \cdots, r\rangle$ be the Weyl group of Lie(G), where $\{s_i\}$ are the simple reflections. We identify the Weyl group W with $\mathrm{Norm}_G(H)/H$. An element

$$\overline{s_i} := x_i(-1)y_i(1)x_i(-1) \tag{3.1}$$

is in $\mathrm{Norm}_G(H)$, which is representative of $s_i \in W = \mathrm{Norm}_G(H)/H$ [6]. For $u \in W$, we denote the length of u by $l(u)$.

We have two kinds of Bruhat decompositions of G as follows:

$$G = \coprod_{u \in W} BuB = \coprod_{u \in W} B_- u B_-.$$

Then, for u, $v \in W$, we define the *double Bruhat cell* $G^{u,v}$ as follows:

$$G^{u,v} := BuB \cap B_- v B_-.$$

This is biregularly isomorphic to a Zariski open subset of an affine space of dimension $r + l(u) + l(v)$ [3, Theorem 1.1].

We also define the *reduced double Bruhat cell* $L^{u,v}$ as follows:

$$L^{u,v} := NuN \cap B_- v B_- \subset G^{u,v}.$$

As is similar to the case $G^{u,v}$, $L^{u,v}$ is biregularly isomorphic to a Zariski open subset of an affine space of dimension $l(u) + l(v)$ [2, Proposition 4.4].

Definition 3.1. Let $u = s_{i_1} \cdots s_{i_n}$ be a reduced expression of $u \in W$ $(i_1, \cdots, i_n \in [1, r])$. Then the finite sequence

$$\mathbf{i} := (i_1, \cdots, i_n)$$

is called a *reduced word* for u.

In this paper, we treat (reduced) Double Bruhat cells of the form $G^{u,e} := BuB \cap B_-$ and $L^{u,e} := NuN \cap B_-$, where $u \in W$ is an element whose reduced word can be written as a left factor of $(1, 2, 3, \cdots, r)^r$:

$$u = (s_1 s_2 \cdots s_r)^{m-1} s_1 \cdots s_{i_n}, \tag{3.2}$$

where $n := l(u)$ is the length of u and $1 \leq i_n \leq r$. Let $\mathbf{i}$ be a reduced word of (u, e):

$$\mathbf{i} = (\underbrace{1, \cdots, r}_{\text{1st cycle}}, \underbrace{1, \cdots, r}_{\text{2nd cycle}}, \cdots, \underbrace{1, \cdots, r}_{m-1 \text{ th cycle}}, \underbrace{1, 2, \cdots, i_n}_{m \text{ th cycle}}). \tag{3.3}$$

Note that $(1, 2, 3, \cdots, r)^r$ is a reduced word of the longest element in W.

3.2 Factorization theorem for type C_r

In this subsection, we shall introduce the isomorphisms between double Bruhat cell $G^{u,e}$ and $H \times (\mathbb{C}^\times)^{l(u)}$, and between $L^{u,e}$ and $(\mathbb{C}^\times)^{l(u)}$. As in the previous section, we consider the case $G := Sp_{2r}(\mathbb{C})$. We set $\mathfrak{g} := \mathrm{Lie}(G)$ with the Cartan decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}$. Let e_i, f_i $(i \in [1, r])$ be the generators of $\mathfrak{n}$, $\mathfrak{n}_-$. For $i \in [1, r]$ and $t \in \mathbb{C}$, we set $x_i(t) := \exp(te_i)$, $y_i := \exp(tf_i)$. Let $\varphi_i : SL_2(\mathbb{C}) \to G$ be the canonical embedding corresponding to simple root α_i. Then we have

$$x_i(t) = \varphi_i \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}, \quad y_i(t) = \varphi_i \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix}. \tag{3.4}$$

For a reduced word $\mathbf{i} = (i_1, \cdots, i_n)$ $(i_1, \cdots, i_n \in [1, r])$, we define a map $x^G_{\mathbf{i}} : H \times \mathbb{C}^n \to G$ as

$$x^G_{\mathbf{i}}(a; t_1, \cdots, t_n) := a \cdot y_{i_1}(t_1) \cdots y_{i_n}(t_n). \tag{3.5}$$

Theorem 3.2. [3, *Theorem* 1.2] *We set $u \in W$ and its reduced word $\boldsymbol{i}$ as in* (3.2) *and* (3.3). *The map $x^G_{\boldsymbol{i}}$ defined above can be restricted to a biregular isomorphism between $H \times (\mathbb{C}^\times)^{l(u)}$ and a Zariski open subset of $G^{u,e}$.*

Next, for $i \in [1, r]$ and $t \in \mathbb{C}^\times$, we define as follows:

$$\alpha_i^\vee(t) := t^{h_i} = \varphi_i \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}, \quad x_{-i}(t) := y_i(t)\alpha_i^\vee(t^{-1}) = \varphi_i \begin{pmatrix} t^{-1} & 0 \\ 1 & t \end{pmatrix}. \tag{3.6}$$

For $\mathbf{i} = (i_1, \cdots, i_n)$ $(i_1, \cdots, i_n \in [1, r])$, we define a map $x^L_{\mathbf{i}} : \mathbb{C}^n \to G$ as

$$x^L_{\mathbf{i}}(t_1, \cdots, t_n) := x_{-i_1}(t_1) \cdots x_{-i_n}(t_n). \tag{3.7}$$

We have the following theorem which is similar to the previous one.

Theorem 3.3. [2, *Proposition* 4.5] *We set $u \in W$ and its reduced word $\boldsymbol{i}$ as in* (3.2) *and* (3.3). *The map $x^L_{\boldsymbol{i}}$ defined above can be restricted to a biregular isomorphism between $(\mathbb{C}^\times)^{l(u)}$ and a Zariski open subset of $L^{u,e}$.*

We define a map $\overline{x}^G_{\mathbf{i}} : H \times (\mathbb{C}^\times)^n \to G^{u,e}$ as

$$\overline{x}^G_{\mathbf{i}}(a; t_1, \cdots, t_n) = a x^L_{\mathbf{i}}(t_1, \cdots, t_n),$$

where $a \in H$ and $(t_1, \cdots, t_n) \in (\mathbb{C}^\times)^n$.

Proposition 3.4. *In the above setting, the map $\overline{x}^G_{\mathbf{i}}$ is a biregular isomorphism between $H \times (\mathbb{C}^\times)^n$ and a Zariski open subset of $G^{u,e}$.*

Proof. In this proof, we use the notation

$$(Y_{1,1}, \cdots, Y_{1,r}, \cdots, Y_{m-1,1}, \cdots, Y_{m-1,r}, Y_{m,1}, \cdots, Y_{m,i_n}) \in (\mathbb{C}^\times)^n$$

for variables instead of $(t_1, \cdots, t_n)$.

We define a map $\phi : H \times (\mathbb{C}^\times)^n \to H \times (\mathbb{C}^\times)^n$ as follows: For

$$\mathbf{Y} := (a; Y_{1,1}, \cdots, Y_{1,r}, \cdots, Y_{m,1}, \cdots, Y_{m,i_n}),$$

we define $\phi(\mathbf{Y}) = (\Phi_a(\mathbf{Y}); \Phi_{1,1}(\mathbf{Y}), \cdots, \Phi_{1,r}(\mathbf{Y}), \cdots, \Phi_{m,1}(\mathbf{Y}), \cdots, \Phi_{m,i_n}(\mathbf{Y}))$ as

$$\Phi_a(\mathbf{Y}) := a \cdot \left(\prod_{j=1}^{m-1} \alpha^\vee_1(Y_{j,1})^{-1} \cdots \alpha^\vee_r(Y_{j,r})^{-1} \right) \cdot \alpha^\vee_1(Y_{m,1})^{-1} \cdots \alpha^\vee_{i_n}(Y_{m,i_n})^{-1},$$

and for $1 \leq s \leq m$,

$$\Phi_{s,l}(\mathbf{Y}) := \begin{cases} \frac{(Y_{s+1,l-1}Y_{s+2,l-1}\cdots Y_{m,l-1})(Y_{s,l+1}Y_{s+1,l+1}\cdots Y_{m,l+1})}{Y_{s,l}(Y_{s+1,l}\cdots Y_{m,l})^2} & \text{if } 1 \leq l < r, \\ \frac{(Y_{s+1,r-1}Y_{s+2,r-1}\cdots Y_{m,r-1})^2}{Y_{s,r}(Y_{s+1,r}\cdots Y_{m,r})^2} & \text{if } l = r, \end{cases} \tag{3.8}$$

where in (3.8), if we see the variables $Y_{\zeta,0}$ $(1 \leq \zeta \leq m)$ and $Y_{m,\xi}$ $(i_n < \xi)$, then we understand $Y_{\zeta,0} = Y_{m,\xi} = 1$. For example, $Y_{s+1,l-1} = 1$ in the case $l = 1$. Note that ϕ is a biregular isomorphism since we can recurrently construct the inverse map $\psi : H \times (\mathbb{C}^\times)^n \to H \times (\mathbb{C}^\times)^n$, $\mathbf{Y} \mapsto (\Psi_a(\mathbf{Y}); \Psi_{1,1}(\mathbf{Y}), \cdots, \Psi_{m,i_n}(\mathbf{Y}))$ of ϕ as follows: The definition (3.8) implies that $\Phi_{m,i_n}(\mathbf{Y}) = \frac{1}{Y_{m,i_n}}$, and hence $Y_{m,i_n} = \frac{1}{\Psi_{m,i_n}(\mathbf{Y})}$. So we set $\Psi_{m,i_n}(\mathbf{Y}) = \frac{1}{Y_{m,i_n}}$. Suppose that we can construct $\Psi_{m,i_n}(\mathbf{Y}), \Psi_{m,i_n-1}(\mathbf{Y}), \cdots, \Psi_{m,1}(\mathbf{Y}), \cdots \Psi_{s+1,r}(\mathbf{Y}), \cdots, \Psi_{s+1,1}(\mathbf{Y}), \Psi_{s,r}(\mathbf{Y}), \cdots, \Psi_{s,l+1}(\mathbf{Y})$. Then we define

$$\Psi_{s,l}(\mathbf{Y}) := \begin{cases} \frac{(\Psi_{s+1,l}(\mathbf{Y})\cdots\Psi_{m,l}(\mathbf{Y}))^2}{Y_{s,l}(\Psi_{s+1,l-1}(\mathbf{Y})\Psi_{s+2,l-1}(\mathbf{Y})\cdots\Psi_{m,l-1}(\mathbf{Y}))(\Psi_{s,l+1}(\mathbf{Y})\cdots\Psi_{m,l+1}(\mathbf{Y}))} & \text{if } 1 \leq l < r, \\ \frac{(\Psi_{s+1,r}(\mathbf{Y})\cdots\Psi_{m,r}(\mathbf{Y}))^2}{Y_{s,r}(\Psi_{s+1,r-1}(\mathbf{Y})\Psi_{s+2,r-1}(\mathbf{Y})\cdots\Psi_{m,r-1}(\mathbf{Y}))^2} & \text{if } l = r. \end{cases}$$

We also define

$$\Psi_a(\mathbf{Y}) := a \cdot \left(\prod_{j=1}^{m-1} \alpha^\vee_1(\Psi_{j,1}(\mathbf{Y})) \cdots \alpha^\vee_r(\Psi_{j,r}(\mathbf{Y})) \right) \cdot \alpha^\vee_1(\Psi_{m,1}(\mathbf{Y})) \cdots \alpha^\vee_{i_n}(\Psi_{m,i_n}(\mathbf{Y})).$$

Then, we get the inverse map ψ of ϕ.

Let us prove

$$\overline{x}^G_{\mathbf{i}}(\mathbf{Y}) = (x^G_{\mathbf{i}} \circ \phi)(\mathbf{Y}),$$

which implies that $\overline{x}^G_{\mathbf{i}} : H \times (\mathbb{C}^\times)^n \to G^{u,e}$ is biregular isomorphism by Theorem 3.2.

First, it is known that

$$\alpha^\vee_i(c)^{-1} y_j(t) = \begin{cases} y_i(c^2 t)\alpha^\vee_i(c)^{-1} & \text{if } i = j, \\ y_j(c^{-1} t)\alpha^\vee_i(c)^{-1} & \text{if } |i-j| = 1 \text{ and } (i,j) \neq (r-1,r), \\ y_j(c^{-2} t)\alpha^\vee_i(c)^{-1} & \text{if } (i,j) = (r-1,r), \\ y_j(t)\alpha^\vee_i(c)^{-1} & \text{otherwise}, \end{cases} \tag{3.9}$$

for $1 \leq i,\ j \leq r$ and $c,\ t \in \mathbb{C}^{\times}$.

On the other hand, it follows from the definition (3.5) of $x_{\mathbf{i}}^{G}$ and (3.8) that

$$
\begin{aligned}
&(x_{\mathbf{i}}^{G} \circ \phi)(\mathbf{Y}) \\
&\quad = a \times \left(\prod_{j=1}^{m-1} \alpha_1^{\vee}(Y_{j,1})^{-1} \cdots \alpha_r^{\vee}(Y_{j,r})^{-1} \right) \cdot \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} \\
&\times y_1(\Phi_{1,1}(\mathbf{Y})) y_2(\Phi_{1,2}(\mathbf{Y})) \cdots y_r(\Phi_{1,r}(\mathbf{Y})) \cdots y_1(\Phi_{m,1}(\mathbf{Y})) \cdots y_{i_n}(\Phi_{m,i_n}(\mathbf{Y})).
\end{aligned} \tag{3.10}
$$

For each s and l $(1 \leq s \leq m,\ 1 \leq l \leq r)$, we can move

$$
\begin{aligned}
&\alpha_l^{\vee}(Y_{s,l})^{-1} \alpha_{l+1}^{\vee}(Y_{s,l+1})^{-1} \cdots \alpha_r^{\vee}(Y_{s,r})^{-1} \\
&\qquad \cdot \left(\prod_{j=s+1}^{m-1} \alpha_1^{\vee}(Y_{j,1})^{-1} \cdots \alpha_r^{\vee}(Y_{j,r})^{-1} \right) \cdot \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1}
\end{aligned}
$$

to the right of $y_l(\Phi_{s,l}(\mathbf{Y}))$ by using the relations (3.9). For example,

$$
\alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} y_l(\Phi_{s,l}(\mathbf{Y})) =
$$

$$
\begin{cases}
y_l \left(\frac{Y_{m,l}^2}{Y_{m,l-1} Y_{m,l+1}} \Phi_{s,l}(\mathbf{Y}) \right) \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} & \text{if } 1 \leq l < r, \\
y_r \left(\frac{Y_{m,r}^2}{Y_{m,r-1}^2} \Phi_{s,r}(\mathbf{Y}) \right) \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} & \text{if } l = r.
\end{cases}
$$

Repeating this argument, in the case $l < r$, we have

$$
\begin{aligned}
&\alpha_l^{\vee}(Y_{s,l})^{-1} \alpha_{l+1}^{\vee}(Y_{s,l+1})^{-1} \cdots \alpha_r^{\vee}(Y_{s,r})^{-1} \\
&\times \left(\prod_{j=s+1}^{m-1} \alpha_1^{\vee}(Y_{j,1})^{-1} \cdots \alpha_r^{\vee}(Y_{j,r})^{-1} \right) \cdot \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} y_l(\Phi_{s,l}(\mathbf{Y})) \\
&= y_l \left(\frac{(Y_{s,l} Y_{s+1,l} \cdots Y_{m-1,l} Y_{m,l})^2}{(Y_{s+1,l-1} \cdots Y_{m-1,l-1} Y_{m,l-1})(Y_{s,l+1} \cdots Y_{m-1,l+1} Y_{m,l+1})} \Phi_{s,l}(\mathbf{Y}) \right) \cdot \alpha_l^{\vee}(Y_{s,l})^{-1} \\
&\times \alpha_{l+1}^{\vee}(Y_{s,l+1})^{-1} \cdots \alpha_r^{\vee}(Y_{s,r})^{-1} \cdot \left(\prod_{j=s+1}^{m-1} \alpha_1^{\vee}(Y_{j,1})^{-1} \cdots \alpha_r^{\vee}(Y_{j,r})^{-1} \right) \cdot \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1}.
\end{aligned}
$$

Note that $\frac{(Y_{s,l} Y_{s+1,l} \cdots Y_{m-1,l} Y_{m,l})^2}{(Y_{s+1,l-1} \cdots Y_{m-1,l-1} Y_{m,l-1})(Y_{s,l+1} \cdots Y_{m-1,l+1} Y_{m,l+1})} \Phi_{s,l}(\mathbf{Y}) = Y_{s,l}$, which implies

$$
\alpha_{l+1}^{\vee}(Y_{s,l+1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} y_l(\Phi_{s,l}(\mathbf{Y})) = y_l(Y_{s,l}) \alpha_{l+1}^{\vee}(Y_{s,l+1})^{-1} \cdots \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1}. \tag{3.11}
$$

In the case $l = r$, we can also verify the relation (3.11) similarly. Thus, by (3.10) and (3.11), we have

$$
\begin{aligned}
(x_{\mathbf{i}}^{G} \circ \phi)(\mathbf{Y}) &= a \cdot y_1(Y_{1,1}) \alpha_1^{\vee}(Y_{1,1})^{-1} \cdots y_r(Y_{1,r}) \alpha_r^{\vee}(Y_{1,r})^{-1} \times \cdots \\
&\quad \times y_1(Y_{m,1}) \alpha_1^{\vee}(Y_{m,1})^{-1} \cdots y_{i_n}(Y_{m,i_n}) \alpha_{i_n}^{\vee}(Y_{m,i_n})^{-1} \\
&= a \cdot x_{-1}(Y_{1,1}) \cdots x_{-r}(Y_{1,r}) \cdots x_{-1}(Y_{m,1}) \cdots x_{-i_n}(Y_{m,i_n}) \\
&= \overline{x}_{\mathbf{i}}^{G}(\mathbf{Y}).
\end{aligned}
$$

□

4 Cluster algebras and generalized minors

For this section, see *e.g.,*[1, 3, 4, 5].

We set $[1,l]:=\{1,2,\cdots,l\}$ and $[-1,-l]:=\{-1,-2,\cdots,-l\}$ for $l\in\mathbb{Z}_{>0}$. For $n,m\in\mathbb{Z}_{>0}$, let $x_1,\cdots,x_n,x_{n+1},\cdots,x_{n+m}$ be variables and $\mathcal{P}$ be a free multiplicative abelian group generated by $x_{n+1},\cdots,x_{n+m}$. We set $\mathbb{Z}\mathcal{P}:=\mathbb{Z}[x_{n+1}^{\pm1},\cdots,x_{n+m}^{\pm1}]$. Let $K:=\{\frac{g}{h}|\ g,\ h\in\mathbb{Z}\mathcal{P},\ h\neq 0\}$ be the field of fractions of $\mathbb{Z}\mathcal{P}$, and $\mathcal{F}:=K(x_1,\cdots,x_n)$ be the field of rational functions.

4.1 Cluster algebras of geometric type

Definition 4.1. Let $B=(b_{ij})$ be an $n\times n$ integer matrix.

(i) B is *skew symmetric* if $b_{ij}=-b_{ji}$ for any $i,\ j\in[1,n]$.

(ii) B is *skew symmetrizable* if there exists a positive integer diagonal matrix D such that DB is skew symmetric.

(iii) B is *sign skew symmetric* if $b_{ij}b_{ji}\leq 0$ for any $i,\ j\in[1,n]$, and if $b_{ij}b_{ji}=0$ then $b_{ij}=b_{ji}=0$.

Note that each skew symmetric matrix is skew symmetrizable, and each skew symmetrizable matrix is sign skew symmetric.

Definition 4.2. We set n-tuple of variables $\mathbf{x}=(x_1,\cdots,x_n)$. Let $\tilde{B}=(b_{ij})_{1\leq i\leq n,\ 1\leq j\leq n+m}$ be $n\times(n+m)$ integer matrix whose principal part $B:=(b_{ij})_{1\leq i,j\leq n}$ is sign skew symmetric. Then a pair $\Sigma=(\mathbf{x},\tilde{B})$ is called a *seed*, $\mathbf{x}$ a cluster and $x_1,\cdots,\ x_n$ cluster variables. For a seed $\Sigma=(\mathbf{x},\tilde{B})$, principal part B of $\tilde{B}$ is called the *exchange matrix*.

Definition 4.3. If B is skew symmetric (resp. skew symmetrizable, sign skew symmetric), we say $\tilde{B}$ is skew symmetric (resp. skew symmetrizable, sign skew symmetric).

Definition 4.4. For a seed $\Sigma=(\mathbf{x},\tilde{B}=(b_{ij}))$, an *adjacent cluster* in direction $k\in[1,n]$ is defined by

$$\mathbf{x}_k=(\mathbf{x}\setminus\{x_k\})\cup\{x'_k\},$$

where x'_k is the new cluster variable defined by the *exchange relation*

$$x_kx'_k=\prod_{1\leq i\leq n+m,\ b_{ki}>0}x_i^{b_{ki}}+\prod_{1\leq i\leq n+m,\ b_{ki}<0}x_i^{-b_{ki}}.$$

Definition 4.5. Let $A=(a_{ij}),\ A'=(a'_{ij})$ be two matrices of the same size. We say that A' is obtained from A by the matrix mutation in direction k, and denote $A'=\mu_k(A)$ if

$$a'_{ij}=\begin{cases}-a_{ij} & \text{if } i=k \text{ or } j=k,\\ a_{ij}+\frac{|a_{ik}|a_{kj}+a_{ik}|a_{kj}|}{2} & \text{otherwise.}\end{cases}$$

For $A,\ A'$, if there exists a finite sequence $(k_1,\cdots,k_s)$, $(k_i\in[1,n])$ such that $A'=\mu_{k_1}\cdots\mu_{k_s}(A)$, we say A is mutation equivalent to A', and denote $A\cong A'$.

Proposition 4.6. [*1*] *For* $k\in[1,n]$, $\mu_k(\mu_k(A))=A$.

Definition 4.7. Let A be a sign skew symmetric matrix. We say A is *totally sign skew symmetric* if any matrix that is mutation equivalent to A is sign skew symmetric. Then a seed $(\mathbf{x}, A)$ is called a totally mutable seed.

Next proposition can be easily verified by the definition of μ_k:

Proposition 4.8. [*1, Proposition*3.6] *Skew symmetrizable matrices are totally sign skew symmetric.*

For a seed $\Sigma = (\mathbf{x}, \tilde{B})$, we say that the seed $\Sigma' = (\mathbf{x}', \tilde{B}')$ is adjacent to Σ if $\mathbf{x}'$ is adjacent to $\mathbf{x}$ in direction k and $\tilde{B}' = \mu_k(\tilde{B})$. Two seeds Σ and Σ_0 are mutation equivalent if one of them can be obtained from another seed by a sequence of pairwise adjacent seeds and we denote $\Sigma \sim \Sigma_0$.

Now let us define the cluster algebra of geometric type.

Definition 4.9. Let $\tilde{B}$ be a skew symmetrizable matrix, and $\Sigma = (\mathbf{x}, \tilde{B})$ a seed. We set $\mathbb{A} := \mathbb{Z}[x_{n+1}, \cdots, x_{n+m}]$. The cluster algebra (of geometric type) $\mathcal{A} = \mathcal{A}(\Sigma)$ over $\mathbb{A}$ associated with seed Σ is defined as the $\mathbb{A}$-subalgebra of $\mathcal{F}$ generated by all cluster variables in all seeds which are mutation equivalent to Σ.

For a seed Σ, we define $\mathbb{Z}\mathcal{P}$-subalgebra $\mathcal{U}(\Sigma)$ of $\mathcal{F}$ by

$$\mathcal{U}(\Sigma) := \mathbb{Z}\mathcal{P}[\mathbf{x}^{\pm 1}] \cap \mathbb{Z}\mathcal{P}[\mathbf{x}_1^{\pm 1}] \cap \cdots \cap \mathbb{Z}\mathcal{P}[\mathbf{x}_n^{\pm 1}].$$

Here, $\mathbb{Z}\mathcal{P}[\mathbf{x}^{\pm 1}]$ is the Laurent polynomial ring in $\mathbf{x}$.

Definition 4.10. Let Σ_0 be a totally mutable seed. We define an *upper cluster algebra* $\overline{\mathcal{A}} = \overline{\mathcal{A}}(\Sigma_0)$ as the intersection of the subalgebras $\mathcal{U}(\Sigma)$ for all seeds $\Sigma \sim \Sigma_0$.

For a totally mutable seed Σ, following the inclusion relation holds [5]:

$$\mathcal{A}(\Sigma) \subset \overline{\mathcal{A}}(\Sigma).$$

4.2 Upper cluster algebra $\overline{\mathcal{A}}(\mathbf{i})$

As in Sect.3, let $G = Sp_{2r}(\mathbb{C})$ be the simple algebraic group of type C_r and W be its Weyl group. We set $u \in W$ and its reduced word $\mathbf{i}$ as in (3.2) and (3.3):

$$u = \underbrace{s_1 s_2 \cdots s_r}_{1\,\text{st cycle}} \underbrace{s_1 \cdots s_r}_{2\,\text{nd cycle}} \cdots \underbrace{s_1 \cdots s_r}_{m-1 \text{ th cycle}} \underbrace{s_1 \cdots s_{i_n}}_{m \text{ th cycle}}, \tag{4.1}$$

$$\mathbf{i} = (\underbrace{1, \cdots, r}_{1\,\text{st cycle}}, \underbrace{1, \cdots, r}_{2\,\text{nd cycle}}, \cdots, \underbrace{1, \cdots, r}_{m-1 \text{ th cycle}}, \underbrace{1, \cdots, i_n}_{m \text{ th cycle}}). \tag{4.2}$$

In this subsection, we define the upper cluster algebra $\overline{\mathcal{A}}(\mathbf{i})$, which satisfies that $\overline{\mathcal{A}}(\mathbf{i}) \otimes \mathbb{C}$ is isomorphic to the coordinate ring $\mathbb{C}[G^{u,e}]$ of the double Bruhat cell [5]. Let i_k ($k \in [1, l(u)]$) be the k-th index of $\mathbf{i}$ from the left.

At first, we define a set e($\mathbf{i}$) as

$$e(\mathbf{i}) := [-1, -r] \cup \{k | \text{There exist some } l > k \text{ such that } i_k = i_l\}.$$

Next, let us define a matrix $\tilde{B} = \tilde{B}(\mathbf{i})$.

Definition 4.11. Let $\tilde{B}(\mathbf{i})$ be an integer matrix with rows labelled by all the indices in $[-1,-r]\cup[1,l(u)]$ and columns labelled by all the indices in $e(\mathbf{i})$. For $k\in[-1,-r]\cup[1,l(u)]$ and $l\in e(\mathbf{i})$, an entry b_{kl} of $\tilde{B}(\mathbf{i})$ is determined as follows:

$$b_{kl}=\begin{cases}-\mathrm{sgn}\,((k-l)\cdot i_p) & \text{if } p=q,\\ -\mathrm{sgn}((k-l)\cdot i_p\cdot a_{|i_k||i_l|}) & \text{if } p<q \text{ and } \mathrm{sgn}(i_p\cdot i_q)(k-l)(k^+-l^+)>0,\\ 0 & \text{otherwise.}\end{cases}$$

Proposition 4.12. [5, *Proposition* 2.6] $\tilde{B}(\boldsymbol{i})$ *is skew symmetrizable.*

By Proposition 4.8, Definition 4.10 and Proposition 4.12, we can construct the upper cluster algebra:

Definition 4.13. We denote this upper cluster algebra by $\overline{\mathcal{A}}(\mathbf{i})$.

4.3 Generalized minors and bilinear form

As in the previous section, we set $G=Sp_{2r}(\mathbb{C})$, $u\in W$ and its reduced word $\mathbf{i}$ as in (4.1) and (4.2). We also set $\bar{\mathcal{A}}(\mathbf{i})_{\mathbb{C}}:=\bar{\mathcal{A}}(\mathbf{i})\otimes\mathbb{C}$ and $\mathcal{F}_{\mathbb{C}}:=\mathcal{F}\otimes\mathbb{C}$. It is known that the coordinate ring $\mathbb{C}[G^{u,e}]$ of the double Bruhat cell is isomorphic to $\bar{\mathcal{A}}(\mathbf{i})_{\mathbb{C}}$ (Theorem 4.15). To describe this isomorphism explicitly, we need generalized minors.

We set $G_0:=N_-HN$, and let $x=[x]_-[x]_0[x]_+$ with $[x]_-\in N_-$, $[x]_0\in H$, $[x]_+\in N$ be the corresponding decomposition.

Definition 4.14. For $i\in[1,r]$ and w, $w'\in W$, the generalized minor $\Delta_{w\Lambda_i,w'\Lambda_i}$ is a regular function on G whose restriction to the open set $wG_0w'^{-1}$ is given by $\Delta_{w\Lambda_i,w'\Lambda_i}(x)=([w^{-1}xw']_0)^{\Lambda_i}$. Here, Λ_i is the i-th fundamental weight. In particular, we write $\Delta_{\Lambda_i}:=\Delta_{\Lambda_i,\Lambda_i}$ and call it *principal minor.*

We set $\mathfrak{g}=\mathrm{Lie}(G)$. Let $\omega:\mathfrak{g}\to\mathfrak{g}$ be the anti involution

$$\omega(e_i)=f_i,\quad \omega(f_i)=e_i,\quad \omega(h)=h,$$

and extend it to G by setting $\omega(x_i(c))=y_i(c)$, $\omega(y_i(c))=x_i(c)$ and $\omega(t)=t$ ($t\in H$). Here, x_i and y_i were defined in Sect.3.2 (3.4).

There exists a $\mathfrak{g}$ (or G)-invariant bilinear form on the finite-dimensional irreducible $\mathfrak{g}$-module $V(\lambda)$ such that

$$\langle au,v\rangle=\langle u,\omega(a)v\rangle,\qquad (u,v\in V(\lambda),\ a\in\mathfrak{g}\text{ (or }G)).$$

For $g\in G$, we have the following simple fact:

$$\Delta_{\Lambda_i}(g)=\langle gu_{\Lambda_i},u_{\Lambda_i}\rangle,$$

where u_{Λ_i} is a properly normalized highest weight vector in $V(\Lambda_i)$. Hence, for $w,w'\in W$, we have

$$\Delta_{w\Lambda_i,w'\Lambda_i}(g)=\Delta_{\Lambda_i}(\overline{w}^{-1}g\overline{w}')=\langle\overline{w}^{-1}g\overline{w}'\cdot u_{\Lambda_i},u_{\Lambda_i}\rangle=\langle g\overline{w}'\cdot u_{\Lambda_i}\,,\,\overline{w}\cdot u_{\Lambda_i}\rangle,\quad(4.3)$$

where $\overline{w}$ is the one we defined in Sect.3.1 (3.1), and note that $\omega(\overline{s}_i^{\pm})=\overline{s}_i^{\mp}$.

4.4 Cluster algebras on Double Bruhat cells of type C

For $k \in [1, l(u)]$, let i_k be the k-th index of $\mathbf{i}$ (4.2) from the left, and we suppose that it belongs to the m' th cycle. We set

$$u_{\leq k} = u_{\leq k}(\mathbf{i}) := \underbrace{s_1 s_2 \cdots s_r}_{1\,\text{st cycle}} \underbrace{s_1 \cdots s_r}_{2\,\text{nd cycle}} \cdots \underbrace{s_1 \cdots s_{i_k}}_{m'\,\text{th cycle}} . \tag{4.4}$$

For $k \in [-1, -r]$, we set $u_{\leq k} := e$ and $i_k := k$. For $k \in [-1, -r] \cup [1,\ l(u)]$, we define

$$\Delta(k; \mathbf{i})(x) := \Delta_{u_{\leq k}\Lambda_{i_k}, \Lambda_{i_k}}(x).$$

Finally, we set

$$F(\mathbf{i}) := \{\Delta(k; \mathbf{i})(x) | k \in [-1, -r] \cup [1,\ l(u)]\}.$$

It is known that the set $F(\mathbf{i})$ is an algebraically independent generating set for the field of rational functions $\mathbb{C}(G^{u,e})$ [3, Theorem 1.12]. Then, we have the following theorem.

Theorem 4.15. [5, *Theorem* 2.10] *The isomorphism of fields* $\varphi : F_{\mathbb{C}} \to \mathbb{C}(G^{u,e})$ *defined by* $\varphi(x_k) = \Delta(k; \boldsymbol{i})$ $(k \in [-1, -r] \cup [1, l(u)])$ *restricts to an isomorphism of algebras* $\bar{\mathcal{A}}(\boldsymbol{i})_{\mathbb{C}} \to \mathbb{C}[G^{u,e}]$.

5 Generalized minors and crystals

In the rest of the paper, we consider the case $G = Sp_{2r}(\mathbb{C})$. Let $u \in W$ be

$$u = (s_1 s_2 \cdots s_r)^{m-1} s_1 \cdots s_{i_n}, \tag{5.1}$$

where $n = l(u)$, $1 \leq i_n \leq r$ and $1 \leq m \leq r$. Let

$$\mathbf{i} = (\underbrace{1, \cdots, r}_{1\,\text{st cycle}}, \underbrace{1, \cdots, r}_{2\,\text{nd cycle}}, \cdots,\ \underbrace{1, \cdots, r}_{m-1\,\text{th cycle}}, \underbrace{1, \cdots, i_n}_{m\,\text{th cycle}}), \tag{5.2}$$

be a reduced word $\mathbf{i}$ for u, that is, $\mathbf{i}$ is the left factor of $(1, 2, 3, \cdots, r)^r$. Let i_k be the k-th index of $\mathbf{i}$ from the left, and belong to m'-th cycle. As we shall show in lemma 5.5, we may assume $i_n = i_k$.

By Theorem 4.15, we can regard $\mathbb{C}[G^{u,e}]$ as an upper cluster algebra and $\{\Delta(k; \mathbf{i})\}$ as its cluster variables. Each $\Delta(k; \mathbf{i})$ is a regular function on $G^{u,e}$. On the other hand, by Proposition 3.4 (resp. Theorem 3.3), we can consider $\Delta(k; \mathbf{i})$ as a function on $H \times (\mathbb{C}^\times)^{l(u)}$ (resp. $(\mathbb{C}^\times)^{l(u)}$). Then we change the variables of $\{\Delta(k; \mathbf{i})\}$ as follows:

Definition 5.1. For $a \in H$ and

$$\mathbf{Y} := (Y_{1,1}, Y_{1,2}, \cdots, Y_{1,r}, Y_{2,1}, Y_{2,2}, \cdots, Y_{2,r},$$
$$\cdots, Y_{m-1,1}, \cdots, Y_{m-1,r}, Y_{m,1}, \cdots, Y_{m,i_n}) \in (\mathbb{C}^\times)^n, \tag{5.3}$$

we set

$$\Delta^G(k; \mathbf{i})(a, \mathbf{Y}) := (\Delta(k; \mathbf{i}) \circ \overline{x}^G_{\mathbf{i}})(a, \mathbf{Y}),$$
$$\Delta^L(k; \mathbf{i})(\mathbf{Y}) := (\Delta(k; \mathbf{i}) \circ x^L_{\mathbf{i}})(\mathbf{Y}).$$

We will describe the function $\Delta^L(k;\mathbf{i})(\mathbf{Y})$ explicitly since $\Delta^G(k;\mathbf{i})(a,\mathbf{Y})$ is immediately obtained from $\Delta^L(k;\mathbf{i})(\mathbf{Y})$ (Proposition 5.3).

Remark 5.2. *If we see the variables $Y_{s,0}$, $Y_{s,r+1}$ $(1\leq s\leq m)$ then we understand*

$$Y_{s,0}=Y_{s,r+1}=1.$$

For example, if $i=1$ then

$$Y_{s,i-1}=1.$$

5.1 Generalized minor $\Delta^G(k;\mathbf{i})(a,\mathbf{Y})$

In this subsection, we shall prove that $\Delta^G(k;\mathbf{i})(a,\mathbf{Y})$ is immediately obtained from $\Delta^L(k;\mathbf{i})(\mathbf{Y})$:

Proposition 5.3. *We set $d:=i_k$. For $a=t^{\sum_i a_i h_i}\in H$ $(t\in\mathbb{C}^\times)$, we have*

$$\Delta^G(k;\boldsymbol{i})(a,\boldsymbol{Y})=\begin{cases} t^{(a_r-a_{m'}-a_{d-r+m'})}\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y}) & \text{if } m'+d>r,\\ t^{(a_{m'+d}-a_{m'})}\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y}) & \text{if } m'+d\leq r.\end{cases}$$

This proposition follows from (2.2) and the following lemma:

Lemma 5.4. *In the above setting, if $m'+d>r$, then we have*

$$\Delta^G(k;\boldsymbol{i})(a,\boldsymbol{Y})=\langle ax^L_{\boldsymbol{i}}(\boldsymbol{Y})(v_1\wedge v_2\wedge\cdots\wedge v_d),\quad v_{m'+1}\wedge\cdots\wedge v_r\wedge v_{\overline{d-r+m'}}\wedge\cdots\wedge v_{\overline{1}}\rangle,$$

$$\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y})=\langle x^L_{\boldsymbol{i}}(\boldsymbol{Y})(v_1\wedge v_2\wedge\cdots\wedge v_d),\quad v_{m'+1}\wedge\cdots\wedge v_r\wedge v_{\overline{d-r+m'}}\wedge\cdots\wedge v_{\overline{1}}\rangle,$$

where $\langle,\rangle$ is the bilinear form we defined in Sect.4.3. In the case $m'+d\leq r$, we have

$$\Delta^G(k;\boldsymbol{i})(a,\boldsymbol{Y})=\langle ax^L_{\boldsymbol{i}}(\boldsymbol{Y})(v_1\wedge v_2\wedge\cdots\wedge v_d),\quad v_{m'+1}\wedge\cdots\wedge v_{m'+d}\rangle,$$

$$\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y})=\langle x^L_{\boldsymbol{i}}(\boldsymbol{Y})(v_1\wedge v_2\wedge\cdots\wedge v_d),\quad v_{m'+1}\wedge\cdots\wedge v_{m'+d}\rangle.$$

Proof. Let us prove this lemma for $\Delta^L(k;\mathbf{i})(\mathbf{Y})$ since the case for $\Delta^G(k;\mathbf{i})(a,\mathbf{Y})$ is proven similarly. Using (4.3) and (4.4), we see that $\Delta^L(k;\mathbf{i})(\mathbf{Y})=\Delta_{u_{\leq k}\Lambda_d,\Lambda_d}(x^L_{\mathbf{i}}(\mathbf{Y}))$ is given as

$$\langle x^L_{\mathbf{i}}(\mathbf{Y})(v_1\wedge v_2\wedge\cdots\wedge v_d),\quad \underbrace{\overline{s_1}\cdots\overline{s_r}}_{1\,\text{st cycle}}\cdots\underbrace{\overline{s_1}\cdots\overline{s_d}}_{m'\,\text{th cycle}}(v_1\wedge v_2\wedge\cdots\wedge v_d)\rangle. \tag{5.4}$$

By (3.1), for $1\leq i\leq r-1$ and $1\leq j\leq r$, we get

$$\overline{s_i}v_j=\begin{cases} v_{i+1} & \text{if } j=i,\\ -v_i & \text{if } j=i+1,\\ v_j & \text{if otherwise,}\end{cases}\qquad \overline{s_i}v_{\overline{j}}=\begin{cases} v_{\overline{i}} & \text{if } j=i+1,\\ -v_{\overline{i+1}} & \text{if } j=i,\\ v_{\overline{j}} & \text{if otherwise,}\end{cases}$$

and we obtain

$$\overline{s_r}v_j=\begin{cases} v_{\overline{r}} & \text{if } j=r,\\ v_j & \text{if } j\neq r,\end{cases}\qquad \overline{s_r}v_{\overline{j}}=\begin{cases} -v_r & \text{if } j=r,\\ v_{\overline{j}} & \text{if } j\neq r.\end{cases}$$

Therefore, if $m' + d \leq r$, then

$$u_{\leq k}(v_1 \wedge \cdots \wedge v_d) = \underbrace{\overline{s_1} \cdots \overline{s_r}}_{1\,\text{st cycle}} \cdots \underbrace{\overline{s_1} \cdots \overline{s_d}}_{m'\,\text{th cycle}} (v_1 \wedge \cdots \wedge v_d) = v_{m'+1} \wedge v_{m'+2} \wedge \cdots \wedge v_{m'+d}. \tag{5.5}$$

If $m' + d > r$, then we get

$$\begin{aligned}
& u_{\leq k}(v_1 \wedge \cdots \wedge v_d) \\
= \;& \underbrace{\overline{s_1} \cdots \overline{s_r}}_{1\,\text{st cycle}} \cdots \underbrace{\overline{s_1} \cdots \overline{s_d}}_{m'\,\text{th cycle}} (v_1 \wedge v_2 \wedge \cdots \wedge v_d) \\
= \;& \underbrace{\overline{s_1} \cdots \overline{s_r}}_{1\,\text{st cycle}} \cdots \underbrace{\overline{s_1} \cdots \overline{s_r}}_{m'-r+d\,\text{th cycle}} (v_{r-d+1} \wedge \cdots \wedge v_r) \\
= \;& \underbrace{\overline{s_1} \cdots \overline{s_r}}_{1\,\text{st cycle}} \cdots \underbrace{\overline{s_1} \cdots \overline{s_r}}_{m'-r+d-1\,\text{th cycle}} (v_{r-d+2} \wedge \cdots \wedge v_r \wedge v_{\overline{1}}) \\
= \;& \underbrace{\overline{s_1} \cdots \overline{s_r}}_{1\,\text{st cycle}} \cdots \underbrace{\overline{s_1} \cdots \overline{s_r}}_{m'-r+d-2\,\text{th cycle}} (v_{r-d+3} \wedge \cdots \wedge v_r \wedge v_{\overline{1}} \wedge (-v_{\overline{2}})) \\
= \;& \cdots \quad = v_{m'+1} \wedge \cdots \wedge v_r \wedge v_{\overline{1}} \wedge (-v_{\overline{2}}) \wedge ((-1)^2 v_{\overline{3}}) \wedge \cdots \wedge ((-1)^{d-r+m'-1} v_{\overline{d-r+m'}}) \\
= \;& v_{m'+1} \wedge \cdots \wedge v_r \wedge v_{\overline{d-r+m'}} \wedge \cdots \wedge v_{\overline{1}}.
\end{aligned} \tag{5.6}$$

Hence, we get our claim by (5.4). □

In the rest of the paper, we will treat $\Delta^L(k; \mathbf{i})(\mathbf{Y})$ only by Proposition 5.3.

5.2 Generalized minor $\Delta^L(k; \mathbf{i})(\mathbf{Y})$

Lemma 5.5. *Let u, $\boldsymbol{i}$ and $\boldsymbol{Y}$ be as in* (5.1), (5.2) *and* (5.3). *Let $i_{n+1} \in [1, r]$ be an index such that $u' := us_{i_{n+1}} \in W$ satisfies $l(u') > l(u)$. We set the reduced word $\boldsymbol{i}'$ for u' as*

$$\boldsymbol{i}' = (\underbrace{1, \cdots, r}_{1\,\text{st cycle}}, \underbrace{1, \cdots, r}_{2\,\text{nd cycle}}, \cdots, \underbrace{1, \cdots, r}_{m-1\,\text{th cycle}}, \underbrace{1, \cdots, i_n}_{m\,\text{th cycle}}, i_{n+1}),$$

and denote $\boldsymbol{Y}' \in (\mathbb{C}^\times)^{n+1}$ by

$$\boldsymbol{Y}' := (Y_{1,1}, \cdots, Y_{1,r}, \cdots, Y_{m-1,1}, \cdots, Y_{m-1,r}, Y_{m,1}, \cdots, Y_{m,i_n}, Y).$$

For an integer k $(1 \leq k \leq n)$, if $d := i_k \neq i_{n+1}$, then $\Delta^L(k; \boldsymbol{i})(\boldsymbol{Y})$ does not depend on Y, so we can regard it as a function on $(\mathbb{C}^\times)^n$. Furthermore, we have

$$\Delta^L(k; \boldsymbol{i})(\boldsymbol{Y}) = \Delta^L(k; \boldsymbol{i}')(\boldsymbol{Y}'). \tag{5.7}$$

Proof. By the definition (3.7) of $x^L_{\mathbf{i}}$, we have

$$x^L_{\mathbf{i}'}(\mathbf{Y}) = x^L_{\mathbf{i}}(\mathbf{Y}) x_{-i_{n+1}}(Y). \tag{5.8}$$

On the other hand, since $f_i^2 = 0$ on $V(\Lambda_1)$, we have $\exp(tf_i) = 1 + tf_i$ $(i = 1, \cdots, r,\ t \in \mathbb{C})$. Hence, by $x_{-i_{n+1}}(Y) := \exp(Yf_{i_{n+1}}) \cdot (Y^{-h_{i_{n+1}}})$ (see (3.6)), we get

$$x_{-i_{n+1}}(Y) v_j = \begin{cases} Y^{-1} v_{i_{n+1}} + v_{i_{n+1}+1} & \text{if } j = i_{n+1}, \\ Y v_{i_{n+1}+1} & \text{if } j = i_{n+1}+1, \\ v_j & \text{otherwise,} \end{cases} \tag{5.9}$$

where in the case $j = i_{n+1}$, we set $v_{r+1} := v_{\overline{r}}$. Thus, if $d < i_{n+1}$, then we have $x_{-i_{n+1}}(Y)(v_1 \wedge \cdots \wedge v_d) = v_1 \wedge \cdots \wedge v_d$. If $d > i_{n+1}$, then we have

$x_{-i_{n+1}}(Y)(v_1 \wedge \cdots \wedge v_d)$

$$\begin{aligned} &= v_1 \wedge \cdots \wedge v_{i_{n+1}-1} \wedge (Y^{-1}v_{i_{n+1}} + v_{i_{n+1}+1}) \wedge Yv_{i_{n+1}+1} \wedge \cdots \wedge v_d \\ &= v_1 \wedge \cdots \wedge v_d. \end{aligned}$$

Since we assume $i_{n+1} \neq d$, we get

$$x_{-i_{n+1}}(Y)(v_1 \wedge \cdots \wedge v_d) = v_1 \wedge \cdots \wedge v_d. \tag{5.10}$$

We can easily see that $u_{\leq k} = u'_{\leq k}(= \underbrace{s_1 \cdots s_r}_{1\,\text{st cycle}} \cdots \underbrace{s_1 \cdots s_d}_{m'\,\text{th cycle}})$. Therefore, it follows from (4.3), (5.8) and (5.10) that

$$\begin{aligned} \Delta^L(k;\mathbf{i}')(\mathbf{Y}') &= \Delta_{u'_{\leq k}\Lambda_d,\Lambda_d}(x^L_{\mathbf{i}'}(\mathbf{Y}')) \\ &= \langle x^L_{\mathbf{i}'}(\mathbf{Y}')(v_1 \wedge v_2 \wedge \cdots \wedge v_d),\ u'_{\leq k}(v_1 \wedge v_2 \wedge \cdots \wedge v_d)\rangle \\ &= \langle x^L_{\mathbf{i}}(\mathbf{Y})(v_1 \wedge v_2 \wedge \cdots \wedge v_d),\ u_{\leq k}(v_1 \wedge v_2 \wedge \cdots \wedge v_d)\rangle = \Delta^L(k;\mathbf{i})(\mathbf{Y}), \end{aligned}$$

which is our desired result. □

By this lemma, when we calculate $\Delta^L(k;\mathbf{i})(\mathbf{Y})$, we may assume that $i_n = i_k$ without loss of generality.

For $1 \leq l \leq m$ and $1 \leq k \leq r$, we set the Laurent monomials

$$\overline{C}(l,k) := \frac{Y_{l,k-1}}{Y_{l,k}}, \quad C(l,k) := \frac{Y_{l,k+1}}{Y_{l+1,k}}. \tag{5.11}$$

Remark 5.6. *In [6], it was defined* $\overline{C}^{(l)}_k := \frac{Y_{r-l,k-1}}{Y_{r-l,k}}$ *and* $C^{(l)}_k := \frac{Y_{r-l,k}}{Y_{r-l+1,k-1}}$, *which coincide with* $\overline{C}(r-l,k)$ *and* $C(r-l,k-1)$ *in* (5.11) *respectively.*

For $1 \leq l \leq r$, we set $|l| = |\overline{l}| = l$. The following theorem is our main result.

Theorem 5.7. *In the above setting, we set* $d := i_k = i_n$ *and*

$$\boldsymbol{Y} := (Y_{1,1}, Y_{1,2}, \cdots, Y_{1,r}, \cdots, Y_{m-1,1}, \cdots, Y_{m-1,r}, Y_{m,1}, \cdots, Y_{m,i_n}) \in (\mathbb{C}^\times)^n.$$

Then we have

$$\begin{aligned} \Delta^L(k;\boldsymbol{i})(\boldsymbol{Y}) = \sum_{(*)} \prod_{i=1}^{d} &\overline{C}(m - l_i^{(1)}, k_i^{(1)}) \cdots \overline{C}(m - l_i^{(\delta_i)}, k_i^{(\delta_i)}) \\ &\cdot C(m - l_i^{(\delta_i+1)}, |k_i^{(\delta_i+1)}| - 1) \cdots C(m - l_i^{(m-m')}, |k_i^{(m-m')}| - 1), \end{aligned}$$

$$l_i^{(s)} := \begin{cases} k_i^{(s)} + s - i - 1 & \text{if } s \leq \delta_i, \\ s - i + r & \text{if } s > \delta_i \end{cases} \qquad (1 \leq i \leq d)$$

where $(*)$ *is the conditions for* $k_i^{(s)}$ $(1 \leq s \leq m-m',\ 1 \leq i \leq d)$: $1 \leq k_1^{(s)} < k_2^{(s)} < \cdots < k_d^{(s)} \leq \overline{1} \quad (1 \leq s \leq m-m')$, $1 \leq k_i^{(1)} \leq \cdots \leq k_i^{(m-m')} \leq m' + i \quad (1 \leq i \leq r-m')$, *and* $1 \leq k_i^{(1)} \leq \cdots \leq k_i^{(m-m')} \leq \overline{1} \quad (r-m'+1 \leq i \leq d)$, *and* δ_i $(i = 1, \cdots, d)$ *are the numbers which satisfy* $1 \leq k_i^{(1)} \leq k_i^{(2)} \leq \cdots \leq k_i^{(\delta_i)} \leq r$, $\overline{r} \leq k_i^{(\delta_i+1)} \leq \cdots \leq k_i^{(m-m')} \leq \overline{1}$.

Example 5.8. *For* rank $r = 3$, $u = s_1s_2s_3s_1s_2s_3s_1s_2$, $k = 5$ *and the reduced word* $\boldsymbol{i} = (-1,-2,-3,-1,-2,-3,-1,-2)$ *for* u*, we have* $m = 3$, $m' = 2$ *and* $d = 2$ *(see* (5.1), (5.2)*). Then, we have* $s = 1$ *and then we write* k_i *for* $k_i^{(s)}$*. Thus, the set of all* (k_1, k_2) *satisfying* $(*)$ *in Theorem 5.7 is*

$$\{(1,2),(1,3),(1,\overline{3}),(1,\overline{2}),(1,\overline{1}),(2,3),(2,\overline{3}),(2,\overline{2}),(2,\overline{1}),(3,\overline{3}),(3,\overline{2}),(3,\overline{1})\}$$

Here, for all (k_1, k_2) *the corresponding monomials are as follows:*

$$\begin{array}{lll}
(1,2) \leftrightarrow \overline{C}(3,1)\overline{C}(3,2) & (1,3) \leftrightarrow \overline{C}(3,1)\overline{C}(2,3) & (1,\overline{3}) \leftrightarrow \overline{C}(3,1)C(1,2) \\
(1,\overline{2}) \leftrightarrow \overline{C}(3,1)C(1,1) & (1,\overline{1}) \leftrightarrow \overline{C}(3,1)C(1,0) & (2,3) \leftrightarrow \overline{C}(2,2)\overline{C}(2,3) \\
(2,\overline{3}) \leftrightarrow \overline{C}(2,2)C(1,2) & (2,\overline{2}) \leftrightarrow \overline{C}(2,2)C(1,1) & (2,\overline{1}) \leftrightarrow \overline{C}(2,2)C(1,0) \\
(3,\overline{3}) \leftrightarrow \overline{C}(1,3)C(1,2) & (3,\overline{2}) \leftrightarrow \overline{C}(1,3)C(1,1) & (3,\overline{1}) \leftrightarrow \overline{C}(1,3)C(1,0)
\end{array}$$

Thus, we obtain:

$$\begin{aligned}
\Delta^L(5;\boldsymbol{i})(\boldsymbol{Y}) &= \overline{C}(3,1)\overline{C}(3,2) + \overline{C}(3,1)\overline{C}(2,3) + \overline{C}(3,1)C(1,2) + \overline{C}(3,1)C(1,1) \\
&\quad + \overline{C}(3,1)C(1,0) + \overline{C}(2,2)\overline{C}(2,3) + \overline{C}(2,2)C(1,2) + \overline{C}(2,2)C(1,1) \\
&\quad + \overline{C}(2,2)C(1,0) + \overline{C}(1,3)C(1,2) + \overline{C}(1,3)C(1,1) + \overline{C}(1,3)C(1,0) \\
&= \frac{1}{Y_{3,2}} + \frac{Y_{2,2}}{Y_{3,1}Y_{2,3}} + \frac{Y_{1,3}}{Y_{3,1}Y_{2,2}} + \frac{Y_{1,2}}{Y_{3,1}Y_{2,1}} + \frac{Y_{1,1}}{Y_{3,1}} + \frac{Y_{2,1}}{Y_{2,3}} + \frac{Y_{2,1}Y_{1,3}}{Y_{2,2}^2} \\
&\quad + 2\frac{Y_{1,2}}{Y_{2,2}} + \frac{Y_{2,1}Y_{1,1}}{Y_{2,2}} + \frac{Y_{1,2}^2}{Y_{2,1}Y_{1,3}} + \frac{Y_{1,1}Y_{1,2}}{Y_{1,3}}.
\end{aligned}$$

Note that since $\overline{C}(1,3)C(1,2) = \overline{C}(2,2)C(1,1) = \frac{Y_{1,2}}{Y_{2,2}}$*, the coefficient of* $\frac{Y_{1,2}}{Y_{2,2}}$ *in the above formula is equal to* 2.

6 The proof of Theorem 5.7

In this section, we shall give the proof of Theorem 5.7.

6.1 The set $X_d(m, m')$ of paths: path descriptions

In this subsection, we shall introduce a set $X_d(m, m')$ of "paths" which correspond to the terms of $\Delta^L(k;\mathbf{i})(\mathbf{Y})$, which we call *path descriptions* of generalized minors. Let m, m' and d be the positive integers as in 5.2. We set $J := \{j, \overline{j} \mid 1 \le j \le r\}$ and for $1 \le l \le r$, set $|l| = |\overline{l}| = l$.

Definition 6.1. Let us define the directed graph (V_d, E_d) as follows: We define the set $V_d = V_d(m)$ of vertices as

$$V_d(m) := \{\mathrm{vt}(m-s; a_1^{(s)}, a_2^{(s)}, \cdots, a_d^{(s)}) | 0 \le s \le m,\ a_i^{(s)} \in J\}.$$

And we define the set $E_d = E_d(m)$ of directed edges as

$$\begin{aligned}
&E_d(m) := \{\mathrm{vt}(m-s; a_1^{(s)}, \cdots, a_d^{(s)}) \to \mathrm{vt}(m-s-1; a_1^{(s+1)}, \cdots, a_d^{(s+1)}) \\
&\mid 0 \le s \le m-1,\ \mathrm{vt}(m-s; a_1^{(s)}, \cdots, a_d^{(s)}),\ \mathrm{vt}(m-s-1; a_1^{(s+1)}, \cdots, a_d^{(s+1)}) \in V_d(m)\}.
\end{aligned}$$

Now, let us define the set of directed paths from $\mathrm{vt}(m;1,2,\cdots,d)$ to $\mathrm{vt}(0;m'+1,m'+2,\cdots,r,\overline{d-r+m'},\overline{d-r+m'-1},\cdots,\overline{2},\overline{1})$ (resp. $\mathrm{vt}(0;m'+1,m'+2,\cdots,m'+d)$) in the case $m'+d>r$ (resp. $m'+d\leq r$) in (V_d,E_d).

Definition 6.2. Let $X_d(m,m')$ be the set of directed paths p

$$p=\mathrm{vt}(m;a_1^{(0)},\cdots,a_d^{(0)})\to\mathrm{vt}(m-1;a_1^{(1)},\cdots,a_d^{(1)})\to\mathrm{vt}(m-2;a_1^{(2)},\cdots,a_d^{(2)})\\ \to\cdots\to\mathrm{vt}(1;a_1^{(m-1)},\cdots,a_d^{(m-1)})\to\mathrm{vt}(0;a_1^{(m)},\cdots,a_d^{(m)}),$$

which satisfy the following conditions: For $0\leq s\leq m$,

(i) $a_\zeta^{(s)}\in J$ $(1\leq\zeta\leq d)$,

(ii) $a_1^{(s)}<a_2^{(s)}<\cdots<a_d^{(s)}$,

(iii) If $a_\zeta^{(s)}\in\{j|1\leq j\leq r-1\}$, then $a_\zeta^{(s+1)}=a_\zeta^{(s)}$ or $a_\zeta^{(s)}+1$. If $a_\zeta^{(s)}=r$, then $a_{\zeta+1}^{(s)}\in\{r,\overline{r},\overline{r-1},\cdots,\overline{1}\}$. If $a_\zeta^{(s)}\in\{\overline{j}|1\leq j\leq r\}$, then $a_\zeta^{(s+1)}\in\{\overline{|a_\zeta^{(s)}|},\overline{|a_\zeta^{(s)}|-1},\cdots,\overline{2},\overline{1}\}$,

(iv) $(a_1^{(0)},a_2^{(0)},\cdots,a_d^{(0)})=(1,2,\cdots,d)$,

$$(a_1^{(m)},\cdots,a_d^{(m)})=\begin{cases}(m'+1,m'+2,\cdots,r,\overline{d-r+m'},\cdots,\overline{2},\overline{1}) & \text{if } m'+d>r,\\ (m'+1,m'+2,\cdots,m'+d) & \text{if } m'+d\leq r,\end{cases}$$

(v) If $a_\zeta^{(s+1)}\in\{\overline{j}|\ 1\leq j\leq r\}$, then $|a_\zeta^{(s+1)}|>|a_{\zeta+1}^{(s)}|$.

Definition 6.3. We say that two vertices $\mathrm{vt}(m-s;a_1^{(s)},\cdots,a_d^{(s)})$ and $\mathrm{vt}(m-s-1;a_1^{(s+1)},\cdots,a_d^{(s+1)})$ are connected if these vertices satisfy the conditions (i), (ii), (iii) and (v) in Definition 6.2.

Define a Laurent monomial associated with each edge of a path in $X_d(m,m')$.

Definition 6.4. Let $p\in X_d(m,m')$ be a path:

$$p=\mathrm{vt}(m;a_1^{(0)},\cdots,a_d^{(0)})\to\mathrm{vt}(m-1;a_1^{(1)},\cdots,a_d^{(1)})\to\mathrm{vt}(m-2;a_1^{(2)},\cdots,a_d^{(2)})\\ \to\cdots\to\mathrm{vt}(1;a_1^{(m-1)},\cdots,a_d^{(m-1)})\to\mathrm{vt}(0;a_1^{(m)},\cdots,a_d^{(m)}).$$

(i) For each $0\leq s\leq m$, we define the *label of the edge* $\mathrm{vt}(m-s;a_1^{(s)},a_2^{(s)},\cdots,a_d^{(s)})\to\mathrm{vt}(m-s-1;a_1^{(s+1)},a_2^{(s+1)},\cdots,a_d^{(s+1)})$ as the Laurent monomial which determined as follows and denote it $Q^{(s)}(p)$: We suppose that $0\leq\delta\leq d$, $1\leq a_1^{(s)}<\cdots<a_\delta^{(s)}\leq r$, and $a_{\delta+1}^{(s)},\cdots,a_d^{(s)}\in\{\overline{i}|\ 1\leq i\leq r\}$. In the case $a_\delta^{(s)}<r$, we set

$$Q^{(s)}(p):=\frac{Y_{m-s,a_1^{(s+1)}-1}}{Y_{m-s,a_1^{(s)}}}\cdots\frac{Y_{m-s,a_\delta^{(s+1)}-1}}{Y_{m-s,a_\delta^{(s)}}}\frac{Y_{m-s,|a_{\delta+1}^{(s)}|}}{Y_{m-s,|a_{\delta+1}^{(s+1)}|-1}}\cdots\frac{Y_{m-s,|a_d^{(s)}|}}{Y_{m-s,|a_d^{(s+1)}|-1}}.$$

In the case $a_\delta^{(s)} = r$, we set

$$Y(a_\delta^{(s+1)}) := \begin{cases} \frac{Y_{m-s,r-1}}{Y_{m-s,r}} & \text{if } a_\delta^{(s+1)} = r, \\ \frac{1}{Y_{m-s,|a_\delta^{(s+1)}|-1}} & \text{if } a_\delta^{(s+1)} \in \{\bar{i} | i = 1, \cdots, r\}, \end{cases}$$

and set

$$Q^{(s)}(p) := \frac{Y_{m-s,a_1^{(s+1)}-1}}{Y_{m-s,a_1^{(s)}}} \cdots \frac{Y_{m-s,a_{\delta-1}^{(s+1)}-1}}{Y_{m-s,a_{\delta-1}^{(s)}}} Y(a_\delta^{(s+1)}) \frac{Y_{m-s,|a_{\delta+1}^{(s)}|}}{Y_{m-s,|a_{\delta+1}^{(s+1)}|-1}} \cdots \frac{Y_{m-s,|a_d^{(s)}|}}{Y_{m-s,|a_d^{(s+1)}|-1}}.$$

(ii) And we define the *label* $Q(p)$ *of the path* p as the product of them:

$$Q(p) := \prod_{s=0}^{m-1} Q^{(s)}(p). \tag{6.1}$$

(iii) For a subpath p'

$$p' = \mathrm{vt}(m-s'; a_1^{(s')}, \cdots, a_d^{(s')}) \to \mathrm{vt}(m-s'-1; a_1^{(s'+1)}, \cdots, a_d^{(s'+1)}) \to \\ \cdots \to \mathrm{vt}(m-s'-1'; a_1^{(s''-1)}, \cdots, a_d^{(s''-1)}) \to \mathrm{vt}(m-s''; a_1^{(s'')}, \cdots, a_d^{(s'')})$$

of p $(0 \le s' < s'' \le m)$, we define the *label of the subpath* p' as

$$Q(p') := \prod_{s=s'}^{s''} Q^{(s)}(p). \tag{6.2}$$

Example 6.5. *Let* $r = m = 3$, $m' = 2$, $d = 2$. *We can describe the paths of* $X_2(3,2)$ *as follows. For simplicity, we denote vertices* $\mathrm{vt}(*;*,*)$ *by* $(*;*,*)$:



Thus, $X_2(3,2)$ *has the following paths*:
$p_1 = (3;1,2) \to (2;1,2) \to (1;2,3) \to (0;3,\bar{1})$,
$p_2 = (3;1,2) \to (2;1,3) \to (1;2,3) \to (0;3,\bar{1})$,

$p_3 = (3;1,2) \to (2;1,3) \to (1;2,\overline{3}) \to (0;3,\overline{1})$,
$p_4 = (3;1,2) \to (2;1,3) \to (1;2,\overline{2}) \to (0;3,\overline{1})$,
$p_5 = (3;1,2) \to (2;1,3) \to (1;2,\overline{1}) \to (0;3,\overline{1})$,
$p_6 = (3;1,2) \to (2;2,3) \to (1;2,3) \to (0;3,\overline{1})$,
$p_7 = (3;1,2) \to (2;2,3) \to (1;2,\overline{3}) \to (0;3,\overline{1})$,
$p_8 = (3;1,2) \to (2;2,3) \to (1;2,\overline{2}) \to (0;3,\overline{1})$,
$p_9 = (3;1,2) \to (2;2,3) \to (1;2,\overline{1}) \to (0;3,\overline{1})$,
$p_{10} = (3;1,2) \to (2;2,3) \to (1;3,\overline{3}) \to (0;3,\overline{1})$,
$p_{11} = (3;1,2) \to (2;2,3) \to (1;3,\overline{2}) \to (0;3,\overline{1})$,
$p_{12} = (3;1,2) \to (2;2,3) \to (1;3,\overline{1}) \to (0;3,\overline{1})$.

Let us calculate the label of the path p_1*. By Definition 6.4* (*iii*)*, the label* $Q^{(0)}(p_1)$ *of the edge* $(3;1,2) \to (2;1,2)$ *is*

$$Q^{(0)}(p_1) = \frac{Y_{3,1-1}}{Y_{3,1}}\frac{Y_{3,2-1}}{Y_{3,2}} = \frac{1}{Y_{3,2}},$$

where we set $Y_{3,0} = 1$ *following Remark 5.2. The labels of the edges* $(2;1,2) \to (1;2,3)$ *and* $(1;2,3) \to (1;3,\overline{1})$ *are as follows:*

$$Q^{(1)}(p_1) = \frac{Y_{2,2-1}}{Y_{2,1}}\frac{Y_{2,3-1}}{Y_{2,2}} = 1, \quad Q^{(2)}(p_1) = \frac{Y_{1,3-1}}{Y_{1,2}}\frac{1}{Y_{1,1-1}} = 1.$$

Therefore, we get $Q(p_1) = \frac{1}{Y_{3,2}}$.

Similarly, we have

$$Q(p_1) = \frac{1}{Y_{3,2}}, \quad Q(p_2) = \frac{Y_{2,2}}{Y_{3,1}Y_{2,3}}, \quad Q(p_3) = \frac{Y_{1,3}}{Y_{3,1}Y_{2,2}}, \quad Q(p_4) = \frac{Y_{1,2}}{Y_{3,1}Y_{2,1}},$$

$$Q(p_5) = \frac{Y_{1,1}}{Y_{3,1}}, \quad Q(p_6) = \frac{Y_{2,1}}{Y_{2,3}}, \quad Q(p_7) = \frac{Y_{2,1}Y_{1,3}}{Y_{2,2}^2}, \quad Q(p_8) = \frac{Y_{1,2}}{Y_{2,2}},$$

$$Q(p_9) = \frac{Y_{2,1}Y_{1,1}}{Y_{2,2}}, \quad Q(p_{10}) = \frac{Y_{1,2}}{Y_{2,2}}, \quad Q(p_{11}) = \frac{Y_{1,2}^2}{Y_{2,1}Y_{1,3}}, \quad Q(p_{12}) = \frac{Y_{1,1}Y_{1,2}}{Y_{1,3}}.$$

Definition 6.6. For each path $p \in X_d(m, m')$

$$\begin{aligned} p = \mathrm{vt}(m; a_1^{(0)}, \cdots, a_d^{(0)}) &\to \mathrm{vt}(m-1; a_1^{(1)}, \cdots, a_d^{(1)}) \to \mathrm{vt}(m-2; a_1^{(2)}, \cdots, a_d^{(2)}) \\ &\to \cdots \to \mathrm{vt}(1; a_1^{(m-1)}, \cdots, a_d^{(m-1)}) \to \mathrm{vt}(0; a_1^{(m)}, \cdots, a_d^{(m)}) \end{aligned}$$

and $i \in \{1, \cdots, d\}$, we call the following sequence

$$a_i^{(0)} \to a_i^{(1)} \to a_i^{(2)} \to \cdots \to a_i^{(m)}$$

an *i-sequence* of p.

We can easily see the following by Definition 6.2 (iii) and (iv): For $1 \leq i \leq d$,

$$i = a_i^{(0)} \leq a_i^{(1)} \leq \cdots \leq a_i^{(m)}, \tag{6.3}$$

in the order (2.1).

6.2 One-to-one correspondence between paths in $X_d(m,m')$ and terms of $\Delta^L(k;\mathbf{i})(\mathbf{Y})$

In this section, we describe the terms in $\Delta^L(k;\mathbf{i})(\mathbf{Y})$ as the paths in $X_d(m,m')$:

Proposition 6.7. *We use the setting and the notations in Sect.5:*

$$u=(s_1s_2\cdots s_r)^{m-1}s_1\cdots s_{i_n},\quad v=e.$$

Then, we have the following:

$$\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y})=\sum_{p\in X_d(m,m')}Q(p).$$

Let us give an overview of the proof of Proposition 6.7. For $1\leq s\leq m$, we define

$$x^{(s)}_{-[1,r]}:=x_{-1}(Y_{s,1})\cdots x_{-r}(Y_{s,r}). \tag{6.4}$$

For $1\leq s\leq m$ and $i_1,\cdots,i_d\in J:=\{i,\overline{i}|1\leq i\leq r\}$, we set

$$(s;i_1,i_2,\cdots,i_d):=\langle x^{(1)}_{-[1,r]}x^{(2)}_{-[1,r]}\cdots x^{(s)}_{-[1,r]}(v_{i_1}\wedge\cdots\wedge v_{i_d}),\quad u_{\leq k}(v_1\wedge\cdots\wedge v_d)\rangle. \tag{6.5}$$

We shall prove $\Delta^L(k;\mathbf{i})(\mathbf{Y})=(m;1,2,\cdots,d)$ in Lemma 6.9 (i). In Lemma 6.9 (ii) and (iii), we shall also prove the recurrence formula for $\{(s;i_1,\cdots,i_d)\}$, which implies that $\Delta^L(k;\mathbf{i})(\mathbf{Y})=(m;1,2,\cdots,d)$ is expressed as a linear combination of $\{(0;j_1,\cdots,j_d)|j_1,\cdots,j_d\in J,\ j_1<\cdots<j_d\}$. Note that if $(j_1,\cdots,j_d)=(m'+1,m'+2,\cdots,r,\overline{d-r+m'},\overline{d-r+m'-1},\cdots\overline{1})$ (resp. $=(m'+1,m'+2,\cdots,m'+d)$), then $(0;j_1,\cdots,j_d)=1$ in the case $m'+d>r$ (resp. $m'+d\leq r$) by (5.5), (5.6) and (6.5). If $(j_1,\cdots,j_d)$ is not as above, then we get $(0;j_1,\cdots,j_d)=0$. As a sequence of this calculation, we obtain Proposition 6.7 .

First, let us see the following lemma. We can verify it in the same way as (5.9).

Lemma 6.8.

$$x_{-i}(Y)v_j=\begin{cases}Y^{-1}v_i+v_{i+1} & \text{if } j=i,\\ Yv_{i+1} & \text{if } j=i+1,\\ v_j & \text{otherwise,}\end{cases}\quad x_{-i}(Y)v_{\overline{j}}=\begin{cases}Y^{-1}v_{\overline{i+1}}+v_{\overline{i}} & \text{if } j=i+1,\\ Yv_{\overline{i}} & \text{if } j=i,\\ v_{\overline{j}} & \text{otherwise,}\end{cases}$$

for all $1\leq i,j\leq r$ *and* $Y\in\mathbb{C}^\times$, *where we set* $v_{r+1}:=v_{\overline{r}}$.

In the next lemma, we set $|l|=|\overline{l}|=l$ for $1\leq l\leq r$.

Lemma 6.9. *(i)* $\Delta^L(k;\boldsymbol{i})(\boldsymbol{Y})=(m;1,\cdots,d)$.

(ii) For $0\leq\delta\leq d$, $1\leq i_1<\cdots<i_\delta\leq r$, $i_{\delta+1},\cdots,i_d\in\{\overline{i}|\ 1\leq i\leq r\}$ *and* $1\leq s\leq m$, *we have the followings:*

In the case $i_\delta<r$,

$$\begin{aligned}&(s;i_1,\cdots,i_\delta,i_{\delta+1},\cdots,i_d)\\ =&\sum_{(j_1,\cdots,j_d)\in V}\frac{Y_{s,j_1-1}}{Y_{s,i_1}}\cdots\frac{Y_{s,j_\delta-1}}{Y_{s,i_\delta}}\frac{Y_{s,|i_{\delta+1}|}}{Y_{s,|j_{\delta+1}|-1}}\cdots\frac{Y_{s,|i_d|}}{Y_{s,|j_d|-1}}\\ &\cdot(s-1;j_1,\cdots,j_\delta,j_{\delta+1},\cdots,j_d),\end{aligned} \tag{6.6}$$

where $(j_1,\cdots,j_d)$ *runs over* $V := \{(j_1,\cdots,j_d)|\ j_1<\cdots<j_\delta,\ j_\zeta = i_\zeta \text{ or } i_\zeta+1\ (1\le\zeta\le\delta),\ j_\zeta\in\{\overline{|i_\zeta|},\ \overline{|i_\zeta|-1},\cdots,\overline{1}\}\ (\delta+1\le\zeta\le d)\}$.

In the case $i_\delta=r$, *we set*

$$Y(j_\delta) := \begin{cases} \frac{Y_{s,r-1}}{Y_{s,r}} & \text{if } j_\delta = r, \\ \frac{1}{Y_{s,|j_\delta|-1}} & \text{if } j_\delta\in\{\overline{i}|i=1,\cdots,r\}. \end{cases}$$

Then we have

$$\begin{aligned}
&(s;i_1,\cdots,i_{\delta-1},r,i_{\delta+1},\cdots,i_d) \\
=& \sum_{(j_1,\cdots,j_d)\in V} \frac{Y_{s,j_1-1}}{Y_{s,i_1}}\cdots\frac{Y_{s,j_{\delta-1}-1}}{Y_{s,i_{\delta-1}}}\cdot Y(j_\delta)\cdot\frac{Y_{s,|i_{\delta+1}|}}{Y_{s,|j_{\delta+1}|-1}}\cdots\frac{Y_{s,|i_d|}}{Y_{s,|j_d|-1}} \qquad (6.7) \\
&\qquad\cdot(s-1;j_1,\cdots,j_{\delta-1},j_\delta,j_{\delta+1},\cdots,j_d),
\end{aligned}$$

where $(j_1,\cdots,j_d)$ *runs over* $V := \{(j_1,\cdots,j_d)|\ j_1<\cdots<j_\delta,\ j_\zeta = i_\zeta \text{ or } i_\zeta+1\ (1\le\zeta\le\delta-1),\ j_\delta\in\{r,\ \overline{r},\ \overline{r-1},\cdots,\ \overline{1}\},\ j_\zeta\in\{\overline{|i_\zeta|},\ \overline{|i_\zeta|-1},\cdots,\overline{1}\}\ (\delta+1\le\zeta\le d)\}$.

(*iii*) *In addition to the assumptions in* (*ii*), *we suppose that* $i_1<\cdots<i_\delta<i_{\delta+1}<\cdots<i_d$ *with the order* (2.1). *If* $i_\delta<r$, *then we can reduce the range* V *of the sum in* (6.6) *to*

$$V' := \{(j_1,\cdots,j_d)\in V|\ |j_l|>|i_{l+1}|\ (\delta+1\le l\le d-1)\}.$$

If $i_\delta=r$, *then we can reduce the range* V *of the sum in* (6.7) *to*

$$V' := \begin{cases} \{(j_1,\cdots,j_d)\in V|\ |j_l|>|i_{l+1}|\ (\delta+1\le l\le d-1)\} & \text{if } j_\delta=r, \\ \{(j_1,\cdots,j_d)\in V|\ |j_l|>|i_{l+1}|\ (\delta\le l\le d-1)\} & \text{if } j_\delta\in\{\overline{i}|1\le i\le r\}. \end{cases}$$

Proof.

(i) By Lemma 6.8, if $i>j$ $(i,j\in\{1,\cdots,r\})$, then we have $x_{-i}(Y)v_j=v_j$. Thus, we get

$$\begin{aligned}
&(m;1,\cdots,d) := \langle x^{(1)}_{-[1,r]}\cdots x^{(m-1)}_{-[1,r]}x^{(m)}_{-[1,r]}(v_1\wedge\cdots\wedge v_d),\ u_{\le k}(v_1\wedge\cdots\wedge v_d)\rangle \\
=\ & \langle x^{(1)}_{-[1,r]}\cdots x^{(m-1)}_{-[1,r]}x_{-1}(Y_{m,1})\cdots x_{-d}(Y_{m,d})(v_1\wedge\cdots\wedge v_d),\ u_{\le k}(v_1\wedge\cdots\wedge v_d)\rangle \\
=\ & \langle x^L_{\mathbf{i}}(\mathbf{Y})(v_1\wedge\cdots\wedge v_d),\quad u_{\le k}(v_1\wedge\cdots\wedge v_d)\rangle = \Delta^L(k;\mathbf{i})(\mathbf{Y}).
\end{aligned}$$

(ii) By Lemma 6.8, for $1\le s\le m$ and $1\le i\le r$, we get

$$x^{(s)}_{-[1,r]}v_i = \begin{cases} \frac{Y_{s,i-1}}{Y_{s,i}}v_i+v_{i+1} & \text{if } 1\le i\le r-1, \\ \frac{Y_{s,r-1}}{Y_{s,r}}v_r+\sum_{j=1}^r\frac{1}{Y_{s,j-1}}v_{\overline{j}} & \text{if } i=r, \end{cases} \qquad (6.8)$$

and

$$x^{(s)}_{-[1,r]}v_{\overline{i}} = \sum_{j=1}^{i}\frac{Y_{s,i}}{Y_{s,j-1}}v_{\overline{j}}, \qquad (6.9)$$

where we set $Y_{s,0}=1$. Since we supposed that $i_{\delta+1},\cdots,i_d\in\{\overline{i}\mid 1\le i\le r\}$, if $i_\delta<r$, then

$$\begin{aligned}
& x^{(1)}_{-[1,r]}\cdots x^{(s-1)}_{-[1,r]}x^{(s)}_{-[1,r]}(v_{i_1}\wedge\cdots\wedge v_{i_\delta}\wedge v_{i_{\delta+1}}\wedge\cdots\wedge v_{i_d})\\
={}& x^{(1)}_{-[1,r]}\cdots x^{(s-1)}_{-[1,r]}(\left(\frac{Y_{s,i_1-1}}{Y_{s,i_1}}v_{i_1}+v_{i_1+1}\right)\wedge\cdots\wedge\left(\frac{Y_{s,i_\delta-1}}{Y_{s,i_\delta}}v_{i_\delta}+v_{i_\delta+1}\right)\\
&\qquad\wedge\left(\sum_{l=1}^{|i_{\delta+1}|}\frac{Y_{s,|i_{\delta+1}|}}{Y_{s,l-1}}v_{\overline{l}}\right)\wedge\cdots\wedge\left(\sum_{l=1}^{|i_d|}\frac{Y_{s,|i_d|}}{Y_{s,l-1}}v_{\overline{l}}\right))\\
={}& \sum_{j_1,\cdots,j_d}(\frac{Y_{s,j_1-1}}{Y_{s,i_1}}\cdots\frac{Y_{s,j_\delta-1}}{Y_{s,i_\delta}}\frac{Y_{s,|i_{\delta+1}|}}{Y_{s,|j_{\delta+1}|-1}}\cdots\frac{Y_{s,|i_d|}}{Y_{s,|j_d|-1}}\\
&\qquad x^{(1)}_{-[1,r]}\cdots x^{(s-1)}_{-[1,r]}(v_{j_1}\wedge\cdots\wedge v_{j_\delta}\wedge v_{j_{\delta+1}}\wedge\cdots\wedge v_{j_d})),
\end{aligned}\tag{6.10}$$

where $(j_1,\cdots,j_d)$ runs over $\{(j_1,\cdots,j_d)\mid j_1<\cdots<j_\delta,\ j_\zeta=i_\zeta$ or $i_\zeta+1$ $(1\le\zeta\le\delta),\ j_\zeta\in\{\overline{|i_\zeta|},\ \overline{|i_\zeta|-1},\cdots,\overline{1}\}$ $(\delta+1\le\zeta\le d)\}$. We remark that

$$\frac{Y_{s,j_\zeta-1}}{Y_{s,i_\zeta}}=\begin{cases}\frac{Y_{s,i_\zeta-1}}{Y_{s,i_\zeta}} & \text{if } j_\zeta=i_\zeta,\\ 1 & \text{if } j_\zeta=i_\zeta+1,\end{cases}$$

for $1\le\zeta\le\delta$.

By pairing both sides in (6.10) with $u_{\le k}(v_1\wedge\cdots\wedge v_d)$, we obtain (6.6). Similarly, we see (6.7) in the case $i_\delta=r$.

(iii) We suppose that $i_\delta<r$. Let $\hat{V}:=V\setminus V'$ be the complementary set. We define the map $\tau:\hat{V}\to\hat{V}$ as follows: Take $(j_1,\cdots,j_\delta,j_{\delta+1},\cdots,j_d)\in\hat{V}$. Let l $(\delta+1\le l\le d-1)$ be the index such that $|j_{\delta+1}|>|i_{\delta+2}|,\cdots,|j_{l-1}|>|i_l|$ and $|j_l|\le|i_{l+1}|$. Since $|j_{l+1}|\le|i_{l+1}|$ by the definition of V, we have $(j_1,\cdots,j_{l+1},j_l,\cdots,j_d)\in\hat{V}$. So, we define $\tau(j_1,\cdots,j_l,j_{l+1},\cdots,j_d):=(j_1,\cdots,j_{l+1},j_l,\cdots,j_d)$. We can easily see that $\tau^2=id_{\hat{V}}$.

In (6.6), $(s-1,j_1,\cdots,j_l,j_{l+1},\cdots,j_d)$ and $(s-1,j_1,\cdots,j_{l+1},j_l,\cdots,j_d)$ have the same coefficient

$$\frac{Y_{s,j_1-1}}{Y_{s,i_1}}\cdots\frac{Y_{s,|i_l|}}{Y_{s,|j_l|-1}}\frac{Y_{s,|i_{l+1}|}}{Y_{s,|j_{l+1}|-1}}\cdots\frac{Y_{s,|i_d|}}{Y_{s,|j_d|-1}}=\frac{Y_{s,j_1-1}}{Y_{s,i_1}}\cdots\frac{Y_{s,|i_l|}}{Y_{s,|j_{l+1}|-1}}\frac{Y_{s,|i_{l+1}|}}{Y_{s,|j_l|-1}}\cdots\frac{Y_{s,|i_d|}}{Y_{s,|j_d|-1}}.$$

Furthermore, by (6.5), we obtain

$$(s-1,j_1,\cdots,j_l,j_{l+1},\cdots,j_d)=-(s-1,j_1,\cdots,j_{l+1},j_l,\cdots,j_d).$$

Therefore, we get $\Sigma_{\hat{V}}=0$ in (6.6), which implies our desired result. We can verify the case $i_\delta=r$ in the same way. □

Proof of Proposition 6.7.

By the definition of V and V' in Lemma 6.9, we see that $(j_1,\cdots,j_d)\in V'$ if and only if the vertices $\mathrm{vt}(s-1;j_1,\cdots,j_d)$ and $\mathrm{vt}(s;i_1,\cdots,i_d)$ are connected (Definition 6.3). Further, the coefficient of $(s-1;j_1,\cdots,j_d)$ in (6.6), (6.7) coincides with the label of the edge between $\mathrm{vt}(s;i_1,\cdots,i_d)$ and $\mathrm{vt}(s-1;j_1,\cdots,j_d)$ (Definition 6.4 (i)). Let us denote it by ${}^{(s)}Q^{i_1,\cdots,i_d}_{j_1,\cdots,j_d}$. Hence, in the case both $i_\delta=r$ and $i_\delta<r$, we get

$$(s;i_1,\cdots,i_d)=\sum_{(j_1,\cdots,j_d)}{}^{(s)}Q^{i_1,\cdots,i_d}_{j_1,\cdots,j_d}\cdot(s-1;j_1,\cdots,j_d),\tag{6.11}$$

where $(j_1,\cdots,j_d)$ runs over the set $\{(j_1,\cdots,j_d)|\ \mathrm{vt}(s-1;j_1,\cdots,j_d)$ and $\mathrm{vt}(s;i_1,\cdots,i_d)$ are connected$\}$. Note that the conditions $|j_l|>|i_{l+1}|$ in V' and $|i_{l+1}|\geq|j_{l+1}|$ in V implies $|j_l|>|j_{l+1}|$, and we get $j_1<j_2<\cdots<j_d$. Using Lemma 6.9 (iii), we obtain the followings in the same way as (6.11):

$$(s-1;j_1,\cdots,j_d)=\sum_{(k_1,\cdots,k_d)} {}^{(s-1)}Q^{j_1,\cdots,j_d}_{k_1,\cdots,k_d}\cdot(s-2;k_1,\cdots,k_d), \tag{6.12}$$

where $(k_1,\cdots,k_d)$ runs over the set $\{(k_1,\cdots,k_d)|\ \mathrm{vt}(s-2;k_1,\cdots,k_d)$ and $\mathrm{vt}(s-1;j_1,\cdots,j_d)$ are connected$\}$, and ${}^{(s-1)}Q^{j_1,\cdots,j_d}_{k_1,\cdots,k_d}$ is the label of the edge between $\mathrm{vt}(s-1;j_1,\cdots,j_d)$ and $\mathrm{vt}(s-2;k_1,\cdots,k_d)$. By (6.11), (6.12), $(s;i_1,\cdots,i_d)$ is a linear combination of $\{(s-2;k_1,\cdots,k_d)\}$, and the coefficient of $(s-2;k_1,\cdots,k_d)$ is as follows:

$$\sum_{(j_1,\cdots,j_d)} {}^{(s)}Q^{i_1,\cdots,i_d}_{j_1,\cdots,j_d}\cdot{}^{(s-1)}\,Q^{j_1,\cdots,j_d}_{k_1,\cdots,k_d}\cdot(s-2;k_1,\cdots,k_d),$$

where $(j_1,\cdots,j_d)$ runs over the set $\{(j_1,\cdots,j_d)|\ \mathrm{vt}(s-1;j_1,\cdots,j_d)$ is connected to the vertices $\mathrm{vt}(s;i_1,\cdots,i_d)$ and $\mathrm{vt}(s-2;k_1,\cdots,k_d)\}$. The coefficient ${}^{(s)}Q^{i_1,\cdots,i_d}_{j_1,\cdots,j_d}\cdot{}^{(s-1)}\,Q^{j_1,\cdots,j_d}_{k_1,\cdots,k_d}$ coincides with the label of subpath (Definition 6.4 (iii))

$$\mathrm{vt}(s;i_1,\cdots,i_d)\to\mathrm{vt}(s-1;j_1,\cdots,j_d)\to\mathrm{vt}(s-2;k_1,\cdots,k_d).$$

Repeating this argument, we see that $(s;i_1,\cdots,i_d)$ is a linear combination of $\{(0;l_1,\cdots,l_d)\}$ $(1\leq l_1<\cdots<l_d\leq\overline{1})$. The coefficient of $(0;l_1,\cdots,l_d)$ is equal to the sum of labels of all subpaths from $\mathrm{vt}(s;i_1,\cdots,i_d)$ to $\mathrm{vt}(0;l_1,\cdots,l_d)$. In the case $m'+d>r$ (resp. $m'+d\leq r$), for $1\leq l_1<\cdots<l_d\leq\overline{1}$, if $(l_1,\cdots,l_d)=(m'+1,m'+2,\cdots,r,\overline{d-r+m'},\cdots,\overline{2},\overline{1})$ (resp. $=(m'+1,m'+2,\cdots,m'+d)$), then we obtain $(0;l_1,\cdots,l_d)=1$ by (5.5), (5.6) and (6.5). If $(l_1,\cdots,l_d)$ is not as above, we obtain $(0;l_1,\cdots,l_d)=0$. Therefore, we see that $(s;i_1,\cdots,i_d)$ is equal to the sum of labels of subpaths from $\mathrm{vt}(s;i_1,\cdots,i_d)$ to $\mathrm{vt}(0;m'+1,m'+2,\cdots,r,\overline{d-r+m'},\cdots,\overline{2},\overline{1})$ (resp. $\mathrm{vt}(0;m'+1,m'+2,\cdots,m'+d)$).

In particular, $\Delta^L(k;\mathbf{i})(\mathbf{Y})=(m;1,2,\cdots,d)$ is equal to the sum of labels of paths in $X_d(m,m')$, which means $\Delta^L(k;\mathbf{i})(\mathbf{Y})=\sum_{p\in X_d(m,m')}Q(p)$. □

Example 6.10. *Let us assume the same setting as Example 5.8, i.e.,* $r=3$, $u=s_1s_2s_3s_1s_2s_3s_1s_2$, $v=e$, $k=5$, $\boldsymbol{i}=(-1,-2,-3,-1,-2,-3,-1,-2)$, $m=3$, $m'=2$ *and* $d=2$. *Therefore, by Example 6.5, we obtain*

$$\begin{aligned}\Delta^L(5;\boldsymbol{i})(\boldsymbol{Y}) &= \frac{1}{Y_{3,2}}+\frac{Y_{2,2}}{Y_{3,1}Y_{2,3}}+\frac{Y_{1,3}}{Y_{3,1}Y_{2,2}}+\frac{Y_{1,2}}{Y_{3,1}Y_{2,1}}+\frac{Y_{1,1}}{Y_{3,1}}+\frac{Y_{2,1}}{Y_{2,3}}\\ &\quad+\frac{Y_{2,1}Y_{1,3}}{Y_{2,2}^2}+2\frac{Y_{1,2}}{Y_{2,2}}+\frac{Y_{2,1}Y_{1,1}}{Y_{2,2}}+\frac{Y_{1,2}^2}{Y_{2,1}Y_{1,3}}+\frac{Y_{1,1}Y_{1,2}}{Y_{1,3}}.\end{aligned}$$

We find that this just coincides with the explicit form of $\Delta^L(5;\boldsymbol{i})(\boldsymbol{Y})$ *in Example 5.8.*

Remark 6.11. *We suppose that* $m'+d\leq r$.

(1) *Definition 6.2 shows that the set* $X_d(m,m')$ *is constituted by paths* p

$$\begin{aligned}p=\mathrm{vt}(m;a_1^{(0)},\cdots,a_d^{(0)})&\to\mathrm{vt}(m-1;a_1^{(1)},\cdots,a_d^{(1)})\to\\ &\cdots\to\mathrm{vt}(1;a_1^{(m-1)},\cdots,a_d^{(m-1)})\to\mathrm{vt}(0;a_1^{(m)},\cdots,a_d^{(m)})\end{aligned}$$

which satisfy the following conditions: For $0\leq s\leq m$, (*i*) $a^{(s)}_{\zeta} \in \{1, \cdots, r\}$ $(1 \leq \zeta \leq d)$,

(*ii*) $a^{(s)}_1 < a^{(s)}_2 < \cdots < a^{(s)}_d$,

(*iii*) $a^{(s+1)}_{\zeta} = a^{(s)}_{\zeta}$ *or* $a^{(s)}_{\zeta} + 1$.

(*iv*) $(a^{(0)}_1, a^{(0)}_2, \cdots, a^{(0)}_d) = (1, 2, \cdots, d)$,
$(a^{(m)}_1, \cdots, a^{(m)}_d) = (m'+1, m'+2, \cdots, m'+d)$.

(2) *By Definition 6.4, the label* $Q^{(s)}(p)$ *of the edge* $\mathrm{vt}(m-s; a^{(s)}_1, a^{(s)}_2, \cdots, a^{(s)}_d) \to \mathrm{vt}(m-s-1; a^{(s+1)}_1, a^{(s+1)}_2, \cdots, a^{(s+1)}_d)$ *is as follows:*

$$Q^{(s)}(p) := \frac{Y_{m-s, a^{(s+1)}_1 - 1}}{Y_{m-s, a^{(s)}_1}} \cdots \frac{Y_{m-s, a^{(s+1)}_d - 1}}{Y_{m-s, a^{(s)}_d}}.$$

(3) *For* $G_A = SL_{r+1}(\mathbb{C})$, *let* B_A *and* $(B_-)_A$ *be two opposite Borel subgroups in* G_A, $N_A \subset B_A$ *and* $(N_-)_A \subset (B_-)_A$ *their unipotent radicals, and* W_A *be the Weyl group of* G_A. *We define a reduced double Bruhat cell as* $L^{u,v}_A := N_A \cdot u \cdot N_A \cap (B_-)_A \cdot v \cdot (B_-)_A$. *We set* $u, v \in W_A$ *and their reduced word* $\boldsymbol{i}_A$ *as*

$$u = \underbrace{s_1 \cdots s_r}_{1\text{ st cycle}} \underbrace{s_1 \cdots s_{r-1}}_{2\text{ nd cycle}} \cdots \underbrace{s_1 \cdots s_{i_n}}_{m\text{ th cycle}}, \quad v = e,$$

$$\boldsymbol{i}_A = (\underbrace{1, \cdots r}_{1\text{ st cycle}}, \underbrace{1 \cdots (r-1)}_{2\text{ nd cycle}} \cdots \underbrace{1, \cdots, i_n}_{m\text{ th cycle}}),$$

where $n = l(u)$ *and* $1 \leq i_n \leq r - m + 1$. *Let* i_k *be the* k*-th index of* $\boldsymbol{i}_A$ *from the left, and belong to* m'*-th cycle. Using Theorem 3.3, we can define* $\Delta^{L_A}(k; \boldsymbol{i}_A)(\boldsymbol{Y}_A) := (\Delta(k; \boldsymbol{i}_A) \circ x^{L_A}_{\boldsymbol{i}_A})(\boldsymbol{Y}_A)$ *in the same way as Definition 5.1, where*

$$\boldsymbol{Y}_A := (Y_{1,1}, Y_{1,2}, \cdots, Y_{1,r}, Y_{2,1}, Y_{2,2}, \cdots, Y_{2,r-1}, \cdots, Y_{m,1}, \cdots, Y_{m,i_n}) \in (\mathbb{C}^{\times})^n,$$

and the map $x^{L_A}_{\boldsymbol{i}_A} : (\mathbb{C}^{\times})^n \overset{\sim}{\to} L^{u,v}_A$ *is defined as in Theorem 3.3.*
Then, we already had seen in [7] that $\Delta^{L_A}(k; \boldsymbol{i}_A)(\boldsymbol{Y}_A) = \sum_{p \in X_d(m,m')} Q(p)$, *where* $X_d(m, m')$ *and the label* $Q = \prod_{s=0}^{m-1} Q^{(s)}(p)$ *is the one we have seen in* (1) *and* (2). *Therefore, it follows from Proposition 6.7 that if* $m' + d \leq r$, *then* $\Delta^L(k; \boldsymbol{i})(\boldsymbol{Y})$ *coincides with* $\Delta^{L_A}(k; \boldsymbol{i}_A)(\boldsymbol{Y}_A)$.

6.3 The properties of paths in $X_d(m, m')$

In this subsection, we shall see some lemmas on $X_d(m, m')$. By Remark 6.11, we suppose that $m' + d > r$. We fix a path $p \in X_d(m, m')$

$$\begin{aligned} p = \mathrm{vt}(m; a^{(0)}_1, \cdots, a^{(0)}_d) \to \cdots \to \mathrm{vt}(2; a^{(m-2)}_1, \cdots, a^{(m-2)}_d) \\ \to \mathrm{vt}(1; a^{(m-1)}_1, \cdots, a^{(m-1)}_d) \to \mathrm{vt}(0; a^{(m)}_1, \cdots, a^{(m)}_d). \quad (6.13) \end{aligned}$$

Lemma 6.12. *For* $p \in X_d(m, m')$ *in* (6.13), i $(1 \le i \le d-1)$ *and* s $(1 \le s \le m)$, *if* $a_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}$, *then we have* $a_{i+1}^{(s-1)} \in \{\overline{j} | 1 \le j \le r\}$ *and*

$$a_i^{(s)} < a_{i+1}^{(s-1)}.$$

Proof.
Using Definition 6.2 (iii) and the assumption $a_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}$, we obtain $a_i^{(s-1)} \in \{r, \overline{r}, \overline{r-1}, \cdots, \overline{1}\}$. Therefore, we also get $a_{i+1}^{(s-1)} \in \{\overline{r}, \overline{r-1}, \cdots, \overline{1}\}$ by Definition 6.2 (ii). Further, it follows from Definition 6.2 (v) that $a_i^{(s)} < a_{i+1}^{(s-1)}$. □

Lemma 6.13. *For* $p \in X_d(m, m')$ *in* (6.13) *and* i $(r - m' + 1 \le i \le d)$, *we obtain*

$$a_i^{(m)} = a_i^{(m-1)} = \cdots = a_i^{(m-i+r-m'+1)} = \overline{d-i+1}. \tag{6.14}$$

Proof.
By Definition 6.2 (iv), we get $a_{r-m'+1}^{(m)} = \overline{d-r+m'}$, and by Lemma 6.12, we also get $\overline{d-r+m'} = a_{r-m'+1}^{(m)} < a_{r-m'+2}^{(m-1)} \le \overline{1}$. Using Lemma 6.12 repeatedly, we obtain $\overline{d-r+m'} = a_{r-m'+1}^{(m)} < a_{r-m'+2}^{(m-1)} < a_{r-m'+3}^{(m-2)} < \cdots < a_d^{(m-d+r-m'+1)} \le \overline{1}$, which means

$$a_i^{(m-i+r-m'+1)} = \overline{d-i+1} \ \ (r - m' + 1 \le i \le d).$$

It follows from (6.3) and Definition 6.2 (iv) that $\overline{d-i+1} = a_i^{(m-i+r-m'+1)} \le a_i^{(m-i+r-m'+2)} \le \cdots \le a_i^{(m-1)} \le a_i^{(m)} = \overline{d-i+1}$, which yields (6.14). □

By this lemma, we get $a_i^{(s)} = \overline{d-i+1}$ for $r-m'+1 \le i$ and $m-i+r-m'+1 \le s \le m$. In the next lemma, we see the properties for $a_i^{(s)}$ $(0 \le s \le m-i+r-m')$.

Lemma 6.14. *For* i $(1 \le i \le d)$ *and* $p \in X_d(m, m')$, *let*

$$a_i^{(0)} \to a_i^{(1)} \to a_i^{(2)} \to \cdots \to a_i^{(m)}$$

be the i*-sequence of the path* p (*Definition* 6.6).

(i) *In the case* $i \le r - m'$,

$$\#\{0 \le s \le m-1 | \ 1 \le a_i^{(s)} \le r, \text{ and } a_i^{(s)} = a_i^{(s+1)}\} = m - m'.$$

(ii) *In the case* $i > r - m'$,

$$\#\{0 \le s \le m-i+r-m' | \ 1 \le a_i^{(s)} \le r \,\text{and}\, a_i^{(s)} = a_i^{(s+1)}\} + \\ \#\{0 \le s \le m-i+r-m' | \ \overline{r} \le a_i^{(s)} \le \overline{1}\} = m - m'.$$

Proof.
(i) In the case $i \le r - m'$, Definition 6.2 (iv) and (6.3) show that

$$i = a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(m)} = m' + i, \quad a_i^{(s+1)} = a_i^{(s)} \text{ or } a_i^{(s)} + 1. \tag{6.15}$$

In particular, we get $1 \le a_i^{(s)} \le r$ for $1 \le s \le m$. By (6.15), we obtain

$$\#\{0 \le s \le m-1 |\ a_i^{(s+1)} = a_i^{(s)} + 1\} = m',$$

which implies $\#\{0 \le s \le m-1 |\ a_i^{(s)} = a_i^{(s+1)}\} = m - m'$.

(ii) In the case $i > r - m'$, by (6.3), we have

$$i = a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(m-i+r-m')} \le \overline{1}.$$

We suppose that

$$i = a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(l)} \le r, \text{ and } \overline{r} \le a_i^{(l+1)} \le \cdots \le a_i^{(m-i+r-m')} \le \overline{1}, \tag{6.16}$$

for some $1 \le l \le m - i + r - m'$. Definition 6.2 (iii) implies that $a_i^{(s+1)} = a_i^{(s)}$ or $a_i^{(s)} + 1$ $(1 \le s \le l-1)$ and $a_i^{(l)} = r$. Therefore,

$$i = a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(l)} = r,\ a_i^{(s+1)} = a_i^{(s)} \text{ or } a_i^{(s)} + 1.$$

So we have $\#\{1 \le s \le l-1 |\ a_i^{(s+1)} = a_i^{(s)}\} = l - (r - i)$ in the same way as (i).

On the other hand, by the assumption $\overline{r} \le a_i^{(l+1)} \le \cdots \le a_i^{(m-i+r-m')} \le \overline{1}$ in (6.16), we clearly see that $\#\{l+1 \le s \le m-i+r-m' |\ \overline{r} \le a_i^{(s)} \le \overline{1}\} = m - i + r - m' - l$. Hence, $\#\{1 \le s \le l-1 |\ a_i^{(s+1)} = a_i^{(s)}\} + \#\{l+1 \le s \le m-i+r-m' |\ \overline{r} \le a_i^{(s)} \le \overline{1}\} = l - (r-i) + m - i + r - m' - l = m - m'$. □

By this lemma, we define $l_i^{(s)} \in \{0, 1, \cdots, m\}$ $(1 \le i \le d,\ 1 \le s \le m - m')$ for the path $p \in X_d(m, m')$ in (6.13) as follows: For $i \le r - m'$, we set $\{l_i^{(s)}\}_{1 \le s \le m-m'}$ $(l_i^{(1)} < \cdots < l_i^{(m-m')})$ as

$$\{l_i^{(1)},\ l_i^{(2)}, \cdots, l_i^{(m-m')}\} := \{s | a_i^{(s)} = a_i^{(s+1)},\ \ 0 \le s \le m-1\}. \tag{6.17}$$

For $i > r - m'$, we set $\{l_i^{(s)}\}_{1 \le s \le m-m'}$ $(l_i^{(1)} < \cdots < l_i^{(m-m')})$ as

$$\begin{aligned} & \{l_i^{(1)},\ l_i^{(2)}, \cdots, l_i^{(m-m')}\} \\ := \ & \{s |\ 1 \le a_i^{(s)} \le r,\ a_i^{(s)} = a_i^{(s+1)},\ 0 \le s \le m - i + r - m'\} \\ \cup \ & \{s |\ \overline{r} \le a_i^{(s)} \le \overline{1},\ 0 \le s \le m - i + r - m'\}. \end{aligned} \tag{6.18}$$

We also set $k_i^{(s)} \in \{j, \overline{j} |\ 1 \le j \le r\}$ $(1 \le i \le d,\ 1 \le s \le m - m')$ as

$$k_i^{(s)} := a_i^{(l_i^{(s)})}. \tag{6.19}$$

Using (6.3) and $l_i^{(1)} < \cdots < l_i^{(m-m')}$, we obtain

$$k_i^{(1)} \le \cdots \le k_i^{(m-m')}. \tag{6.20}$$

For $1 \le i \le d$, let us define δ_i $(0 \le \delta_i \le m - m')$ as

$$1 \le k_i^{(1)} \le \cdots \le k_i^{(\delta_i)} \le r < \overline{r} \le k_i^{(\delta_i+1)} \le \cdots \le k_i^{(m-m')} \le \overline{1}, \tag{6.21}$$

which is uniquely determined from $\{k_i^{(s)}\}_{s=1,\cdots,m-m'}$.

Lemma 6.15. *(i) For $1 \leq i \leq d$,*

$$l_i^{(s)} = \begin{cases} k_i^{(s)} + s - i - 1 & \text{if } k_i^{(s)} \in \{j | 1 \leq j \leq r\}, \\ s - i + r & \text{if } k_i^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}. \end{cases}$$

(ii) For $1 \leq s \leq m - m'$ and $1 \leq i \leq d - 1$, if $k_i^{(s)} \in \{j | 1 \leq j \leq r\}$, then

$$k_i^{(s)} < k_{i+1}^{(s)}, \quad l_i^{(s)} \leq l_{i+1}^{(s)}.$$

For $1 \leq i \leq d - 1$, if $k_i^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}$, then

$$k_i^{(s)} < k_{i+1}^{(s)}, \quad l_i^{(s)} = l_{i+1}^{(s)} + 1.$$

Proof.

(i) We suppose that $k_i^{(s)} \in \{j | 1 \leq j \leq r\}$. The definition of $l_i^{(s)}$ in (6.17) means that the path p has the following i-sequence (Definition 6.6):

$$\begin{aligned}
&a_i^{(0)} = i,\ a_i^{(1)} = i + 1,\ a_i^{(2)} = i + 2, \cdots, a_i^{(l_i^{(1)})} = i + l_i^{(1)}, \\
&a_i^{(l_i^{(1)}+1)} = i + l_i^{(1)},\ a_i^{(l_i^{(1)}+2)} = i + l_i^{(1)} + 1, \cdots, a_i^{(l_i^{(2)})} = i + l_i^{(2)} - 1, \\
&a_i^{(l_i^{(2)}+1)} = i + l_i^{(2)} - 1,\ a_i^{(l_i^{(2)}+2)} = i + l_i^{(2)}, \cdots, a_i^{(l_i^{(3)})} = i + l_i^{(3)} - 2, \\
&\qquad\qquad\qquad\qquad\vdots \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad (6.22) \\
&a_i^{(l_i^{(s-1)}+1)} = i + l_i^{(s-1)} - s + 2,\ a_i^{(l_i^{(s-1)}+2)} = i + l_i^{(s-1)} - s + 3, \cdots, a_i^{(l_i^{(s)})} = i + l_i^{(s)} - s + 1, \\
&a_i^{(l_i^{(s)}+1)} = i + l_i^{(s)} - s + 1,\ a_i^{(l_i^{(s)}+2)} = i + l_i^{(s)} - s + 2, \cdots.
\end{aligned}$$

Hence we have

$$k_i^{(s)} = a_i^{(l_i^{(s)})} = i + l_i^{(s)} - s + 1, \tag{6.23}$$

which implies $l_i^{(s)} = k_i^{(s)} + s - i - 1$.

Next, we suppose that $a_i^{(l_i^{(s)})} = k_i^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}$. Using (6.3), we get $a_i^{(l_i^{(s)})} \leq a_i^{(l_i^{(s)}+1)} \leq \cdots \leq a_i^{(m-i+r-m')}$ and $a_i^{(\zeta)} \in \{\overline{j} | 1 \leq j \leq r\}$ $(l_i^{(s)} \leq \zeta \leq m - i + r - m')$. Thus, by the definition (6.18) of $l_i^{(s)}$, we obtain $l_i^{(m-m')} = m - i + r - m'$, $l_i^{(m-m'-1)} = m - i + r - m' - 1, l_i^{(m-m'-2)} = m - i + r - m' - 2, \cdots, l_i^{(\xi)} = \xi - i + r$ $(s \leq \xi \leq m - m')$. In particular, we get

$$l_i^{(s)} = s - i + r. \tag{6.24}$$

(ii) We suppose that $k_i^{(s)} \in \{j | 1 \leq j \leq r\}$. If $k_{i+1}^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}$, then we obtain $k_i^{(s)} < k_{i+1}^{(s)}$ in the order (2.1), and it follows $l_i^{(s)} \leq l_{i+1}^{(s)}$ from (i). So we may assume that $k_{i+1}^{(s)} \in \{j | 1 \leq j \leq r\}$.

By Definition 6.2 (ii) and the definition (6.18) of $l_{i+1}^{(s)}$, we have $a_i^{(l_{i+1}^{(s)}+1)} < a_{i+1}^{(l_{i+1}^{(s)}+1)} = a_{i+1}^{(l_{i+1}^{(s)})} = k_{i+1}^{(s)} \leq r$. Therefore, the inequality (6.3) implies

$$i = a_i^{(0)} \leq a_i^{(1)} \leq \cdots \leq a_i^{(l_{i+1}^{(s)})} \leq a_i^{(l_{i+1}^{(s)}+1)} < r, \tag{6.25}$$

$$a_i^{(\zeta)} = a_i^{(\zeta-1)} \text{ or } a_i^{(\zeta-1)} + 1 \ (1 \le \zeta \le l_{i+1}^{(s)} + 1).$$

We obtain

$$l_{i+1}^{(s)} + 1 - s \ge \#\{\zeta | a_i^{(\zeta)} = a_i^{(\zeta-1)} + 1,\ 1 \le \zeta \le l_{i+1}^{(s)} + 1\}, \tag{6.26}$$

otherwise, it follows from (6.25) and (i) that $a_i^{(l_{i+1}^{(s)}+1)} > i + l_{i+1}^{(s)} + 1 - s = k_{i+1} - 1 = a_{i+1}^{(l_{i+1}^{(s)})} - 1$, and hence $a_i^{(l_{i+1}^{(s)}+1)} \ge a_{i+1}^{(l_{i+1}^{(s)})}$, which contradicts Definition 6.2 (ii).

The inequality (6.26) means that

$$s \le \#\{\zeta | a_i^{(\zeta)} = a_i^{(\zeta-1)},\ 1 \le \zeta \le l_{i+1}^{(s)} + 1\}. \tag{6.27}$$

On the other hand, the definition of $l_i^{(s)}$ implies $a_i^{(l_i^{(s)}+1)} = a_i^{(l_i^{(s)})} = k_i^{(s)} \in \{j | 1 \le j \le r\}$. The inequality (6.3) shows

$$i = a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(l_i^{(s)})} = a_i^{(l_i^{(s)}+1)} = k_i^{(s)},$$

$$a_i^{(\zeta)} = a_i^{(\zeta-1)} \text{ or } a_i^{(\zeta-1)} + 1 \ (1 \le \zeta \le l_i^{(s)} + 1),$$

and

$$s = \#\{\zeta | a_i^{(\zeta)} = a_i^{(\zeta-1)},\ 1 \le \zeta \le l_i^{(s)} + 1\}. \tag{6.28}$$

Since $a_i^{(l_i^{(s)})} = a_i^{(l_i^{(s)}+1)}$, the equation (6.28) means

$$s - 1 = \#\{\zeta | a_i^{(\zeta)} = a_i^{(\zeta-1)},\ 1 \le \zeta \le l_i^{(s)}\}. \tag{6.29}$$

Thus, by (6.27) and (6.29), we have $l_i^{(s)} < l_{i+1}^{(s)} + 1$, and hence $l_i^{(s)} \le l_{i+1}^{(s)}$, which yields $k_i^{(s)} < k_{i+1}^{(s)}$ since $k_i^{(s)} = i + l_i^{(s)} - s + 1 < i + l_{i+1}^{(s)} - s + 2 = (i+1) + l_{i+1}^{(s)} - s + 1 = k_{i+1}^{(s)}$.

Next, we suppose that $a_i^{(l_i^{(s)})} = k_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}$. As we have seen in Lemma 6.12, we obtain $a_{i+1}^{(l_i^{(s)}-1)} \in \{\overline{j} | 1 \le j \le r\}$. Since $a_{i+1}^{(l_i^{(s)}-1)} \le a_{i+1}^{(l_i^{(s)})} \le \cdots \le a_{i+1}^{(m-(i+1)+r-m')}$, we get $a_{i+1}^{(\zeta)} \in \{\overline{j} | 1 \le j \le r\}$ $(l_i^{(s)} - 1 \le \zeta \le m - (i+1) + r - m')$ and $l_{i+1}^{(m-m')} = m - (i+1) + r - m'$, $l_{i+1}^{(m-m'-1)} = m - (i+1) + r - m' - 1, \cdots, l_{i+1}^{(\xi)} = \xi - (i+1) + r, \cdots$ $(s \le \xi \le m - m')$ by the definition (6.18) of $l_{i+1}^{(\xi)}$. In particular, we get

$$l_{i+1}^{(s)} = s - (i+1) + r.$$

Therefore, it follows from (6.24) that $l_i^{(s)} = l_{i+1}^{(s)} + 1$. Further, $k_i^{(s)} = a_i^{(l_i^{(s)})} < a_{i+1}^{(l_i^{(s)}-1)} = a_{i+1}^{(l_{i+1}^{(s)})} = k_{i+1}^{(s)}$ by Lemma 6.12. □

6.4 The proof of Theorem 5.7

In this subsection, we shall prove Theorem 5.7. First, we see the following lemma. Let us recall the definition (5.11) of C and $\overline{C}$.

Lemma 6.16. *For $p \in X_d(m,m')$ in* (6.13)*, we set $l_i^{(s)}$, $k_i^{(s)}$ and δ_i as in* (6.17)*,* (6.18)*,* (6.19) *and* (6.21)*. Then we have*

$$\begin{aligned} Q(p) = \prod_{i=1}^{d} &\overline{C}(m - l_i^{(1)}, k_i^{(1)}) \cdots \overline{C}(m - l_i^{(\delta_i)}, k_i^{(\delta_i)}) \\ &\cdot C(m - l_i^{(\delta_i+1)}, |k_i^{(\delta_i+1)}| - 1) \cdots C(m - l_i^{(m-m')}, |k_i^{(m-m')}| - 1). \end{aligned} \tag{6.30}$$

Proof.

At first, we get $a_i^{(l_i^{(\delta_i+1)}-1)} \le r$, otherwise, we have $\overline{r} \le a_i^{(l_i^{(\delta_i+1)}-1)}$ and hence $l_i^{(\delta_i)} = l_i^{(\delta_i+1)} - 1$ and $\overline{r} \le a_i^{(l^{(\delta_i)})} = k_i^{(\delta_i)}$ by the definition (6.18) of $l_i^{(s)}$, which contradicts the assumption of δ_i. Further, we get $a_i^{(l_i^{(\delta_i+1)}-1)} = r$ by $a_i^{(l_i^{(\delta_i+1)})} = k_i^{(\delta_i+1)} \in \{\overline{j} | 1 \le j \le r\}$ and Definition 6.2 (iii). Hence we obtain

$$1 \le a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(l_i^{(\delta_i+1)}-1)} = r < \overline{r} \le a_i^{(l_i^{(\delta_i+1)})} \le \cdots \le a_i^{(m)} \le \overline{1}. \tag{6.31}$$

Next, for $0 \le s \le m-1$ and $1 \le i \le d$, we set the label $Q(a_i^{(s)} \to a_i^{(s+1)})$ as follows:

$$Q(a_i^{(s)} \to a_i^{(s+1)}) := \begin{cases} \dfrac{Y_{m-s, a_i^{(s+1)}-1}}{Y_{m-s, a_i^{(s)}}} & \text{if } 1 \le a_i^{(s)} \le a_i^{(s+1)} \le r, \\ \dfrac{1}{Y_{m-s, |a_i^{(s+1)}|-1}} & \text{if } a_i^{(s)} = r \text{ and } \overline{r} \le a_i^{(s+1)} \le \overline{1}, \\ \dfrac{Y_{m-s, |a_i^{(s)}|}}{Y_{m-s, |a_i^{(s+1)}|-1}} & \text{if } \overline{r} \le a_i^{(s)} \le a_i^{(s+1)} \le \overline{1}, \end{cases} \tag{6.32}$$

which means that the label $Q^{(s)}(p)$ of the edge $\mathrm{vt}(m-s; a_1^{(s)}, \cdots, a_d^{(s)}) \to \mathrm{vt}(m-s-1; a_1^{(s+1)}, \cdots, a_d^{(s+1)})$ is as follows (see Definition 6.4 (i)):

$$Q^{(s)}(p) = \prod_{i=1}^{d} Q(a_i^{(s)} \to a_i^{(s+1)}).$$

Therefore, we get

$$Q(p) = \prod_{s=0}^{m-1} \prod_{i=1}^{d} Q(a_i^{(s)} \to a_i^{(s+1)}),$$

which is obtained from Definition 6.4 (ii). To calculate $\prod_{s=0}^{m-1} Q(a_i^{(s)} \to a_i^{(s+1)})$ for $1 \le i \le d$, let us divide the range of product $\prod_{s=0}^{m-1}$ as follows:

$$\prod_{s=0}^{l_i^{(\delta_i+1)}-2}, \qquad \prod_{s=l_i^{(\delta_i+1)}-1}^{l_i^{(m-m')}} \qquad \text{and} \qquad \prod_{s=l_i^{(m-m')}+1}^{m-1},$$

where in the case $\delta_i = m - m'$, we set

$$l_i^{(m-m'+1)} := l_i^{(m-m')} + 2. \tag{6.33}$$

First, let us consider the first range of the product. For $0 \leq s \leq l_i^{(\delta_i+1)} - 2$, using (6.31) and (6.32), we get

$$Q(a_i^{(s)} \to a_i^{(s+1)}) = \begin{cases} \frac{Y_{m-s,a_i^{(s)}-1}}{Y_{m-s,a_i^{(s)}}} = \overline{C}(m-s, a_i^{(s)}) & \text{if } a_i^{(s+1)} = a_i^{(s)}, \\ 1 & \text{if } a_i^{(s+1)} = a_i^{(s)} + 1, \end{cases}$$

which means

$$\prod_{s=0}^{l_i^{(\delta_i+1)}-2} \left(Q(a_i^{(s)} \to a_i^{(s+1)})\right) = \prod_{\zeta=1}^{\delta_i} \overline{C}(m-\zeta, k_i^{(\zeta)}), \tag{6.34}$$

by (6.18) and $k_i^{(\zeta)} := a_i^{(l_i^{(\zeta)})}$.

Next, we consider the second range of the product. If $r - m' \geq i$ then $r \geq m' + i = a_i^{(m)} \geq \cdots \geq a_i^{(1)} \geq a_i^{(0)}$, which implies $\delta_i = m - m'$, and $\prod_{s=l_i^{(\delta_i+1)}-1}^{l_i^{(m-m')}} \left(Q(a_i^{(s)} \to a_i^{(s+1)})\right) = 1$ by (6.33). So we consider the case $r - m' < i$. For $s = l_i^{(\delta_i+1)} - 1$, we get $Q(a_i^{(s)} \to a_i^{(s+1)}) = \frac{1}{Y_{m-s,|a_i^{(s+1)}|-1}}$, and for $l_i^{(\delta_i+1)} \leq s \leq l_i^{(m-m')}$, $Q(a_i^{(s)} \to a_i^{(s+1)}) = \frac{Y_{m-s,|a_i^{(s)}|}}{Y_{m-s,|a_i^{(s+1)}|-1}}$ by (6.31) and (6.32). Thus, we obtain

$$\begin{aligned} &\prod_{s=l_i^{(\delta_i+1)}-1}^{l_i^{(m-m')}} \left(Q(a_i^{(s)} \to a_i^{(s+1)})\right) \\ = \;& \left(\frac{1}{Y_{m-l_i^{(\delta_i+1)}+1,|a_i^{(l_i^{(\delta_i+1)})}|-1}}\right) \cdot \prod_{s=l_i^{(\delta_i+1)}}^{l_i^{(m-m')}} \frac{Y_{m-s,|a_i^{(s)}|}}{Y_{m-s,|a_i^{(s+1)}|-1}} \\ = \;& \left(\prod_{s=l_i^{(\delta_i+1)}}^{l_i^{(m-m')}} \frac{Y_{m-s,|a_i^{(s)}|}}{Y_{m-s+1,|a_i^{(s)}|-1}}\right) \cdot \frac{1}{Y_{m-l_i^{(m-m')},|a_i^{(l_i^{(m-m')}+1)}|-1}} \\ = \;& \left(\prod_{\zeta=\delta_i+1}^{m-m'} C(m - l_i^{(\zeta)}, |k_i^{\zeta}| - 1)\right) \cdot \frac{1}{Y_{m'+i-r,d-i}}, \end{aligned} \tag{6.35}$$

where for the third equality, we used $l_i^{(m-m')} = m - m' - i + r$ (Lemma 6.15) and $|a_i^{(m-m'-i+r+1)}| = d - i + 1$ (Lemma 6.13).

Finally, we consider the last range of the product. Using Lemma 6.13, Lemma 6.15 and (6.17), we obtain

$$\prod_{s=l_i^{(m-m')}+1}^{m-1} \left(Q(a_i^{(s)} \to a_i^{(s+1)})\right) = \begin{cases} \prod_{s=m-m'-i+r+1}^{m-1} \frac{Y_{m-s,d-i+1}}{Y_{m-s,d-i}} & \text{if } r - m' < i, \\ 1 & \text{if } r - m' \geq i. \end{cases} \tag{6.36}$$

By (6.34), (6.35) and (6.36), to prove (6.30), we need to show that

$$\prod_{i=r-m'+1}^{d} \left(\frac{1}{Y_{m'+i-r,d-i}} \prod_{s=m-m'-i+r+1}^{m-1} \frac{Y_{m-s,d-i+1}}{Y_{m-s,d-i}}\right) = 1. \tag{6.37}$$

We set

$$A := \prod_{i=r-m'+1}^{d} \left(\frac{1}{Y_{m'+i-r,d-i}} \right), \text{ and } \quad B := \prod_{i=r-m'+1}^{d} \left(\prod_{s=m-m'-i+r+1}^{m-1} \frac{Y_{m-s,d-i+1}}{Y_{m-s,d-i}} \right).$$

We obtain the followings:

$$A = \prod_{i=r-m'+1}^{d} \left(\frac{1}{Y_{m'+i-r,d-i}} \right) = \prod_{i=r-m'+1}^{d-1} \left(\frac{1}{Y_{m'+i-r,d-i}} \right) = \prod_{k=1}^{m'+d-r-1} \left(\frac{1}{Y_{k,d-r+m'-k}} \right),$$

and

$$\begin{aligned} B &= \prod_{i=r-m'+1}^{d} \left(\prod_{s=m-m'-i+r+1}^{m-1} \frac{Y_{m-s,d-i+1}}{Y_{m-s,d-i}} \right) = \prod_{i=r-m'+1}^{d} \left(\prod_{s=1}^{m'+i-r-1} \frac{Y_{s,d-i+1}}{Y_{s,d-i}} \right) \\ &= \prod_{s=1}^{m'+d-r-1} \left(\frac{Y_{s,d-r+m'-s}}{Y_{s,d-r+m'-s-1}} \frac{Y_{s,d-r+m'-s-1}}{Y_{s,d-r+m'-s-2}} \frac{Y_{s,d-r+m'-s-2}}{Y_{s,d-r+m'-s-3}} \cdots \frac{Y_{s,1}}{Y_{s,0}} \right) \\ &= \prod_{s=1}^{m'+d-r-1} Y_{s,d-r+m'-s}, \end{aligned}$$

where note that $Y_{s,0} = 1$ (see Remark 5.2). Thus we have $A \cdot B = 1$, which implies (6.37). □

Let us prove the main theorem.

Proof of Theorem 5.7.

Using Lemma 6.16, we see that $Q(p)$ ($p \in X_d(m, m')$) is described as (6.30) with $\{k_i^{(s)}\}_{1 \le i \le d, 1 \le s \le m-m'}$ which satisfy the conditions in Lemma 6.15 (ii), that is, $1 \le k_1^{(s)} < k_2^{(s)} < \cdots < k_d^{(s)} \le \overline{1}$. If $m' + i \le r$, then $a_i^{(0)} \le a_i^{(1)} \le \cdots \le a_i^{(m)} = m' + i$, which means that $1 \le k_i^{(1)} \le k_i^{(2)} \le \cdots \le k_i^{(m-m')} \le m' + i$ for $1 \le i \le r - m'$. For $r - m' + 1 \le i \le d$, the inequality (6.3) implies $1 \le k_i^{(1)} \le k_i^{(2)} \le \cdots \le k_i^{(m-m')} \le \overline{1}$. Thus, $\{k_i^{(s)}\}$ satisfies the conditions $(*)$ in Theorem 5.7.

Conversely, let $\{K_i^{(s)}\}_{1 \le i \le d, 1 \le s \le m-m'}$ the set of numbers which satisfies the conditions $(*)$ in Theorem 5.7:

$$1 \le K_1^{(s)} < K_2^{(s)} < \cdots < K_d^{(s)} \le \overline{1} \quad (1 \le s \le m - m'), \tag{6.38}$$

$$1 \le K_i^{(1)} \le \cdots \le K_i^{(m-m')} \le m' + i \quad (1 \le i \le r - m'), \tag{6.39}$$

and

$$1 \le K_i^{(1)} \le \cdots \le K_i^{(m-m')} \le \overline{1} \quad (r - m' + 1 \le i \le d). \tag{6.40}$$

We need to show that there exists a path $p \in X_d(m, m')$ such that

$$\begin{aligned} Q(p) = \prod_{i=1}^{d} & \overline{C}(m - L_i^{(1)}, K_i^{(1)}) \cdots \overline{C}(m - L_i^{(\delta_i)}, K_i^{(\delta_i)}) \\ & \cdot C(m - L_i^{(\delta_i+1)}, |K_i^{(\delta_i+1)}| - 1) \cdots C(m - L_i^{(m-m')}, |K_i^{(m-m')}| - 1), \end{aligned} \tag{6.41}$$

where δ_i $(1 \le \delta_i \le m - m')$ are the numbers which satisfy $1 \le K_i^{(1)} \le \cdots \le K_i^{(\delta_i)} \le r < \overline{r} \le K_i^{(\delta_i+1)} \le \cdots \le K_i^{(m-m')} \le \overline{1}$, and

$$L_i^{(s)} := \begin{cases} K_i^{(s)} + s - i - 1 & \text{if } K_i^{(s)} \in \{j | 1 \le j \le r\}, \\ s - i + r & \text{if } K_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}, \end{cases}$$

for $1 \le s \le m - m'$ and $1 \le i \le d$. Since we supposed $K_i^{(s)} < K_{i+1}^{(s)}$, we can easily verify

$$L_i^{(s)} \le L_{i+1}^{(s)} \quad \text{if } K_i^{(s)} \in \{j | 1 \le j \le r\}, \tag{6.42}$$

and

$$L_i^{(s)} = L_{i+1}^{(s)} + 1 \quad \text{if } K_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}. \tag{6.43}$$

We claim that $0 \le L_i^{(s)} \le m-1$. By the condition (6.38), we get $i \le K_i^{(s)}$. So it is clear that $0 \le L_i^{(s)}$. For $1 \le i \le r-m'$ and $1 \le s \le m-m'$, it follows from the condition (6.39) that $L_i^{(s)} = K_i^{(s)}+s-i-1 \le m'+i+s-i-1 = m'+s-1 \le m-1$. For $r - m' < i$, we get $L_i^{(s)} \le r - i + s < m' + s \le m$. Therefore, we have $0 \le L_i^{(s)} \le m - 1$ for all $1 \le i \le d$ and $1 \le s \le m - m'$.

Note that if $K_i^{(s)} \in \{\overline{j} | 1 \le j \le r\}$, then $\overline{r} \le K_i^{(s)} < K_{i+1}^{(s)} \le \overline{1}$ and hence

$$L_i^{(s+1)} = L_i^{(s)} + 1. \tag{6.44}$$

We define a path $p = \mathrm{vt}(m; a_1^{(0)}, \cdots, a_d^{(0)}) \to \cdots \to \mathrm{vt}(0; a_1^{(m)}, \cdots, a_d^{(m)}) \in X_d(m, m')$ as follows: For i $(1 \le i \le r-m')$, we define the i-sequence (Definition 6.6) of p as

$$\begin{gathered}
a_i^{(0)} = i,\ a_i^{(1)} = i+1,\ a_i^{(2)} = i+2, \cdots, a_i^{(L_i^{(1)})} = i + L_i^{(1)}, \\
a_i^{(L_i^{(1)}+1)} = i + L_i^{(1)},\ a_i^{(L_i^{(1)}+2)} = i + L_i^{(1)} + 1, \cdots, a_i^{(L_i^{(2)})} = i + L_i^{(2)} - 1, \\
a_i^{(L_i^{(2)}+1)} = i + L_i^{(2)} - 1,\ a_i^{(L_i^{(2)}+2)} = i + L_i^{(2)}, \cdots, a_i^{(L_i^{(3)})} = i + L_i^{(3)} - 2,
\end{gathered}$$

$$\vdots \tag{6.45}$$

$$\begin{gathered}
a_i^{(L_i^{(m-m'-1)}+1)} = i + L_i^{(m-m'-1)} - m + m' + 2,\ \cdots, a_i^{(L_i^{(m-m')})} = \\
i + L_i^{(m-m')} - m + m' + 1, \\
a_i^{(L_i^{(m-m')}+1)} = i+L_i^{(m-m')}-m+m'+1,\ a_i^{(L_i^{(m-m')}+2)} = i+L_i^{(m-m')}-m+m'+2, \\
a_i^{(L_i^{(m-m')}+3)} = i + L_i^{(m-m')} - m + m' + 3, \cdots, a_i^{(m)} = m' + i.
\end{gathered}$$

For i $(r - m' + 1 \le i \le d)$, we define the i-sequence of p as

$$\begin{gathered}
a_i^{(0)} = i,\ a_i^{(1)} = i+1,\ a_i^{(2)} = i+2, \cdots, a_i^{(L_i^{(1)})} = i + L_i^{(1)}, \\
a_i^{(L_i^{(1)}+1)} = i + L_i^{(1)},\ a_i^{(L_i^{(1)}+2)} = i + L_i^{(1)} + 1, \cdots, a_i^{(L_i^{(2)})} = i + L_i^{(2)} - 1, \\
a_i^{(L_i^{(2)}+1)} = i + L_i^{(2)} - 1,\ a_i^{(L_i^{(2)}+2)} = i + L_i^{(2)}, \cdots, a_i^{(L_i^{(3)})} = i + L_i^{(3)} - 2,
\end{gathered}$$

$$\vdots \tag{6.46}$$

$$
\begin{aligned}
&a_i^{(L_i^{(\delta_i-1)}+1)} = i + L_i^{(\delta_i-1)} - \delta_i + 2,\ \cdots, a_i^{(L_i^{(\delta_i)})} = i + L_i^{(\delta_i)} - \delta_i + 1,\\
&a_i^{(L_i^{(\delta_i)}+1)} = i + L_i^{(\delta_i)} - \delta_i + 1,\ a_i^{(L_i^{(\delta_i)}+2)} = i + L_i^{(\delta_i)} - \delta_i + 2,\\
&a_i^{(L_i^{(\delta_i)}+3)} = i + L_i^{(\delta_i)} - \delta_i + 3, \cdots,\ a_i^{(L_i^{(\delta_i+1)}-1)} = r,\\
&a_i^{(L_i^{(\delta_i+1)})} = K_i^{(\delta_i+1)},\ a_i^{(L_i^{(\delta_i+2)})} = K_i^{(\delta_i+2)}, \cdots,\ a_i^{(L_i^{(m-m')})} = K_i^{(m-m')},\\
&a_i^{(L_i^{(m-m')}+1)} = a_i^{(L_i^{(m-m')}+2)} = \cdots = a_i^{(m)} = \overline{d-i+1}.
\end{aligned}
$$

It is easy to see that $a_i^{(L_i^{(s)})} = K_i^{(s)}$ $(1 \leq s \leq m - m')$ by the above lists. Clearly, the path p satisfies Definition 6.2 (iii) and (iv). For $1 \leq s \leq L_i^{(\delta_i+1)} - 1$, we obtain $a_i^{(s)} < a_{i+1}^{(s)}$ by (6.42). For $\delta_i + 1 \leq s \leq m - m'$, we obtain $a_i^{(L_i^{(s)})} < a_{i+1}^{(L_i^{(s)})}$ since $a_i^{(L_i^{(s)})} = K_i^{(s)} < K_{i+1}^{(s)} = a_{i+1}^{(L_{i+1}^{(s)})} = a_{i+1}^{(L_i^{(s)}-1)} \leq a_{i+1}^{(L_i^{(s)})}$ by (6.38) and (6.43). For $L_i^{(m-m')} + 1 \leq s \leq m$, we obtain $a_i^{(s)} = \overline{d-i+1}$, and then we get $a_{i+1}^{(s)} = \overline{d-i}$ since $L_{i+1}^{(m-m')} = L_i^{(m-m')} - 1 < L_i^{(m-m')} \leq s$, which means $a_i^{(s)} < a_{i+1}^{(s)}$. Therefore, $a_i^{(s)} < a_{i+1}^{(s)}$ for all $1 \leq i \leq d-1$ and $1 \leq s \leq m - m'$, which means the path p satisfies Definition 6.2 (ii).

Finally, for $a_i^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}$, we need to verify $a_i^{(s)} < a_{i+1}^{(s-1)}$. The definition (6.45), (6.46) of i-sequence of p shows that either $s = L_i^{(\zeta)}$ for some ζ $(\delta_i \leq \zeta \leq m - m')$ or $L_i^{(m-m')} < s$. In the case $s = L_i^{(\zeta)}$, using (6.38) and (6.43), we see that $a_i^{(s)} = a_i^{(L_i^{(\zeta)})} = K_i^{(\zeta)} < K_{i+1}^{(\zeta)} = a_{i+1}^{(L_{i+1}^{(\zeta)})} = a_{i+1}^{(L_i^{(\zeta)}-1)} = a_{i+1}^{(s-1)}$. In the case $L_i^{(m-m')} < s$, we obtain $a_i^{(s)} = \overline{d-i+1} < \overline{d-i} = a_{i+1}^{(s-1)}$ since $L_{i+1}^{(m-m')} = L_i^{(m-m')} - 1 < L_i^{(m-m')} < s$. Therefore, we have $a_i^{(s)} < a_{i+1}^{(s-1)}$ for $a_i^{(s)} \in \{\overline{j} | 1 \leq j \leq r\}$, which means the path p satisfies Definition 6.2 (v).

Hence p is well-defined, and (6.41) is follows from Lemma 6.16, and Theorem 5.7 follows from Proposition 6.7. $\square$

References

[1] Michael Gekhtman, Michael Shapiro, Alek Vainshtein, Cluster Algebras and Poisson Geometry, AMS (2010).

[2] Arkady Berenstein and Andrei Zelevinsky, Tensor product multiplicities, canonical bases and totally positive varieties, Invent. Math. 143 No. 1, 77–128 (2001).

[3] Sergey Fomin, and Andrei Zelevinsky, Double Bruhat cells and totally positivity. J. Amer. Math. Soc., Vol.12, No 2, 335–380 (1998).

[4] Cluster algebras I: Foundations, Sergey Fomin, and Andrei Zelevinsky, J. Amer. Math. Soc., Vol.15, No.2, 497–529 (2002)

[5] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky, Cluster algebras III : Upper bounds and double Bruhat cells. Duke Mathematical Journal, Vol. 126 No1, 1–52 (2005).

[6] Nakashima T., Decorations on Geometric Crystals and Monomial Realizations of Crystal Bases for Classical Groups, Journal of Algebra, Vol.399, 1, Feb. 712–769 (2014).

[7] Kanakubo Y. and Nakashima T., Cluster variables on double Bruhat cells of type (u, e) and monomial realizations of crystal bases of type A, (preprint) arXiv:1409.8622.